

Journal of Advanced Medical and Dental Sciences Research

@Society of Scientific Research and Studies

NLM ID: 101716117

Journal home page: www.jamdsr.com

doi: 10.21276/jamdsr

Index Copernicus value = 85.10

(e) ISSN Online: 2321-9599;

(p) ISSN Print: 2348-6805

Original Research

Correlation of platelet count and periodontal condition in patients with thrombocytopenia

Dr. Pushpendra Yadav¹, Dr. Rajnish Agarwal²

¹Consultant at Shubham General and Dental Hospital Bikaner, Rajasthan, India;

² Professor Department of Prosthodontics Surendra Dental College & Research Institute Sri Ganganagar Rajasthan

ABSTRACT:

Introduction: Periodontitis is an infectious disease of the supporting structures of the teeth and is triggered by complex, anaerobic, gram-negative specific periodontal bacteria. Thrombocytopenia is a hematological disease associated with decrease in the platelet count in the peripheral blood. **Material and methods:** A case control study was carried out in the Department of Oral medicine and diagnosis at Surendra dental college & Hospital Shri ganganagar. The study was divided into two groups, Group 1 and Group 2 comprising of 15 cases with thrombocytopenia and 15 cases without thrombocytopenia respectively. Both groups were examined for periodontal condition as well as blood tests was done to determine the platelet count. **Results:** In the present study, mean platelet count was higher in Group II as compared to Group I but the difference between the two groups was found to be statistically significant. The severity of periodontitis was greater in Group 1 than that of Group 2 and majority of the patients in Group 1 revealed severe periodontitis. **Conclusion:** Plaque control and motivation are of prime importance in thrombocytopenic subjects so as to prevent gingival hemorrhage and periodontal disease.

Keywords: Thrombocytopenia, Platelet count, Periodontitis.

Received: 23/08/2020

Modified: 26/09/2020

Accepted: 28/09/2020

Corresponding Author: Dr. Rajnish Agarwal, Professor Department of Prosthodontics Surendra Dental College & Research Institute Sri Ganganagar Rajasthan

This article may be cited as: Yadav P, Agarwal R. Correlation of platelet count and periodontal condition in patients with thrombocytopenia. J Adv Med Dent Scie Res 2020;8(10):62-64.

INTRODUCTION:

Periodontitis is an extensive infectious disease of the periodontium which leads to gingival inflammation and bleeding.^{1,2} The main cause of periodontal diseases include a group of specific periodontal pathogens, the host's immunologic and inflammatory response to the bacteria which result in periodontal destruction.³ The aim of the present study was to evaluate the relation between the platelet count and periodontal condition in patients with thrombocytopenia. Periodontal disease constitutes a substantial challenge to the patients as well as oral healthcare professionals. In periodontal infection, biofilm formation, inflammation and attachment loss occur.

The normal count of the platelets varies from 150,000 to 450,000/mm³ whereas in thrombocytopenia, platelet

count falls below 150,000/mm³.^{4,5} Thrombocytopenia manifests clinically as mucocutaneous petechiae, ecchymosis, and hemorrhagic blisters with spontaneous bleeding.⁶ Platelets are small, anucleate cells that travel as resting discoid fragments in the circulation. Platelets are formed and released into the blood stream from megakaryocytes that reside in the bone marrow. Thrombocytopenia is a hematological disease associated with decrease in the platelet count in the peripheral blood and this occurs due to reduced production of platelets by megakaryocytes in the bone marrow, increased destruction of platelets in the circulation and by abnormal sequestration of platelets in the spleen, frequently leads to episodes of hemorrhages. They play an important role in primary hemostasis by the formation of platelet plug as they adhere, spread and

aggregate at the site of vessel injury which in turn prevents the hemorrhage. Apart from these, platelets also possess non-hemostatic properties like angiogenesis, tissue repair, inflammation and metastasis of cancer.⁷ Various studies have suggested that reduced platelet count might be considered as a risk factor for bleeding.⁸⁻¹⁰ For patients with thrombocytopenia, decreased platelet activation, platelet aggregation and thrombopoiesis which are reflected by the reduced presence of reticulated platelets and thus associated with bleeding phenotype.¹¹

MATERIAL AND METHOD

A case control study was carried out in the Department of Oral Medicine and diagnosis Surendra Dental College and Hospital, Sri ganganagar. The study was divided into two groups, Group 1 and Group 2 comprising of 15 cases with thrombocytopenia and 15 cases without thrombocytopenia respectively. In Group 1, all the patients with platelet count less than $150 \times 10^3/\mu\text{L}$ on peripheral blood film were included in our study. On the basis of platelet count, the cases were divided into four grades¹²

Grade I- $75-150 \times 10^3/\mu\text{L}$

Grade II- $50-75 \times 10^3/\mu\text{L}$

Grade III- $25-50 \times 10^3/\mu\text{L}$

Grade IV- $<25 \times 10^3/\mu\text{L}$

In Group 2, all the patients with platelet count in the range of $150 \times 10^3/\mu\text{L}$ - $450 \times 10^3/\mu\text{L}$ were taken in this study. Individuals with a known history of jaundice, blood/blood component transfusions or major surgery, medication, systemic diseases, any other hematological disorders were excluded from the study. Both the groups were examined for periodontal condition for all

the teeth except third molars through plaque index (PI), gingival index (GI), pocket depth (PD) and clinical attachment level (CAL). PD is the distance from the gingival margin to the base of the gingival sulcus or periodontal pocket. CAL is the distance from the cementoenamel junction to the base of the sulcus or periodontal pocket. Healthy subjects displayed periodontal pocket depth <3 mm, clinical attachment loss <3 mm, with no clinical sign of gingivitis but periodontitis patients exhibited ≥ 4 mm of pocket depth and clinical attachment loss ≥ 3 mm in more than 30% sites assessed in the oral cavity. On the basis of severity, periodontitis were classified as mild, moderate and severe.¹³

RESULTS:

In the present study, the mean age of the patients in Group 1 and Group 2 were 51.28 years and 47.32 years respectively. Majority of the patients in Group 1 and Group 2 were males and females respectively. The mean platelet count in Group 1 and Group 2 were $58.62 \pm 5.37 \times 10^3/\mu\text{L}$ and $236.82 \pm 54.32 \times 10^3/\mu\text{L}$ respectively. The p-value was found to be statistically significant between the two groups [Table 1].

The percentage of mild, moderate and severe periodontitis in Group 1 were 6.66%, 26.66% and 66.66% respectively. In Group 2, 6.66% and 93.33% of the subjects showed mild and moderate periodontitis respectively. The severity of periodontitis was higher in Group 1 as compared to Group 2 [Table 2]. In the present study, most of the patients in Group 1 presented with Grade I (33.33%) and Grade III (33.33%) followed by Grade II (26.67%) and Grade IV (6.66%) thrombocytopenia [Table 3].

TABLE 1 Comparison of platelet count between two groups

GROUP	N	PLATELET COUNT MEAN \pm SD	P VALUE
GROUP 1	15	58.62 ± 5.37	
GROUP 2	15	236.82 ± 54.32	.002

GROUP 1 WITH THROMBOCYTOPENIA, GROUP 2 WITHOUT THROMBOCYTOPENIA

TABLE 2 Severity of periodontitis in both the groups

GROUP	N	Mild Periodontitis	Moderate Periodontitis	Severe Periodontitis
GROUP 1	15	1	4	10
GROUP 2	15	1	14	0

TABLE 3 Severity of thrombocytopenia in Group 1

GRADE	NO OF CASE	PERCENTAGE
GRADE 1($75-150 \times 10^3/\mu\text{L}$)	5	33.33 %
GRADE 2($50-75 \times 10^3/\mu\text{L}$)	4	26.67 %
GRADE 3($25-50 \times 10^3/\mu\text{L}$)	5	33.33 %
GRADE 4($<25 \times 10^3/\mu\text{L}$)	1	6.67 %
TOTAL	15	100

DISCUSSION:

Periodontitis is an infectious disease of the supporting structures of the teeth and is triggered by complex, anaerobic, gram-negative specific periodontal bacteria. Periodontal destruction may occur due to the action of toxic products released from the specific pathogenic bacteria along with the host responses initiated against the periodontal bacteria and their products.¹⁴ Thrombocytopenia occur due to either a decreased production of platelets or suppression of the normal blood cells, frequently leads to episodes of hemorrhages.¹⁵ The present study determined the relation between the platelet count and periodontal condition in patients with thrombocytopenia.

In the present study, mean platelet count was higher in Group II as compared to Group I but the difference between the two groups was found to be statistically significant. These results were similar to the case report done by Sangwan et al in 2013. The severity of periodontitis was greater in Group 1 than that of Group 2 and majority of the patients in Group 1 revealed severe periodontitis. In the current study, majority of the patients in Group 1 presented with Grade I and Grade III followed by Grade II and Grade IV thrombocytopenia.

Various studies have revealed that the mean platelet count in subjects with chronic periodontitis was higher as compared to the healthy group but the difference between the two groups was found to be statistically non-significant.^{16,17} However, a study carried out by Gomes et al in 2015 determined that platelet count did not have direct influence on the gingivitis and degree of immune thrombocytopenic purpura, as well as did not have direct relation to the severity of periodontal attachment loss.¹⁸

Platelets are essential for primary hemostasis and endothelial repair, but also play a key role in atherogenesis and thrombus formation. In thrombocytopenia, reduced platelet count, platelet activation and platelet aggregation are associated with bleeding risk. Patients with thrombocytopenia lack effective oral hygiene due to increased risk of bleeding as these subjects might have higher incidence of periodontal condition.

CONCLUSION:

The present study showed that the decrease in the platelet was related to the severe periodontitis in patients with thrombocytopenia. Gingival health is of great importance in these patients as they are more prone to gingival bleeding. Plaque control and motivation are of prime importance in thrombocytopenic subjects so as to prevent gingival hemorrhage and periodontal disease.

REFERENCES:

- Loos BG, Craandijk J, Hoek FJ, Wertheim-van Dillen P, M.van der Velden U. Elevation of systemic markers related to cardiovascular diseases in the peripheral blood of periodontitis patients. *J Periodontol* 2000; 71:1528–1534.
- Nicu EA, Vander Velden U, Nieuwland R, Everts V and Loos BG. Elevated platelet and leukocyte response to oral bacteria in periodontitis. *J Thromb Haemost* 2009; 7 (1): 162–170.
- Listgarten MA. A perspective on periodontal diagnosis. *J Clin Periodontol* 1986; 13:175–81.
- Vinhol PJ, Hvas AM and Nybo M. An overview of platelet indices and methods for evaluating platelet function in thrombocytopenic patients. *Eur J Haematol* 2014; 92:367–376.
- Khammissa RAG, Fourie J, Masilana A, Lawrence S, Lemmer J and Feller L. Oral manifestations of thrombocytopenia. *Saudi Dent J* 2018; 30: 19–25.
- Kashiwagi H and Tomiyama Y. Pathophysiology and management of primary immune thrombocytopenia. *Int J Hematol* 2013; 98:24–33.
- Golebiewska EM and Poole AW. Platelet secretion: from haemostasis to wound healing and beyond. *Blood Rev* 2015; 29:153–162.
- Slichter SJ, Kaufman RM, Assmann SF, McCullough J, Triulzi DJ, Strauss RG et al. Dose of prophylactic platelet transfusions and prevention of hemorrhage. *N Engl J Med* 2010; 362:600–613.
- Uhl L, Assmann SF, Hamza TH, Harrison RW, Gernsheimer T and Slichter SJ. Laboratory predictors of bleeding, and effects of platelet and RBC transfusions on bleeding outcomes, in the PLADO Trial. *Blood* 2017; 130:1247–1258.
- Gaydos LA, Freireich EJ and Mantel N. The quantitative relation between platelet count and hemorrhage in patients with acute leukaemia. *N Engl J Med* 1962; 266:905–909.
- Vinhol PJ. The role of platelets in bleeding in patients with thrombocytopenia and hematological disease. *Clin Chem Lab Med* 2019; 57(12):1808–1817.
- Paramjit E, Rao R, Sudhamani S, Roplekar P, Shaffi Z and Roy S. Spectrum of thrombocytopenia: A clinicopathological study with review of the literature. *Muller J Med Sci Res* 2016; 7(2):121–124.
- Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol* 2007; 78:1387–1399.
- López R, Loos BG, Baelum V. Hematological features in adolescents with Periodontitis. *Clin Oral Invest* 2012; 16:1209–1216.
- Ayers KMS, Colquhoun ANK. Leukemia in children. Part I: Orofacial complications and side-effects of treatment. *N Z Dent J* 2000; 96(424):60–65.
- Abdulaziz Al-Rasheed. Elevation of white blood cells and platelet counts in patients having chronic periodontitis. *Saudi Dent J* 2012; 24(1): 17–21.
- Nicu EA, Velden UVD, Nieuwland R, Everts V, Lo O SBG. Elevated platelet and leukocyte response to oral bacteria in periodontitis. *J Thrombosis Haemostasis*, 7: 162–170.
- Gomes de RJ, Lima E, Pastrana AV and de Souza. Study of the relation between platelet count and periodontal condition in patients with immune thrombocytopenic purpura. *J Res Dent* 2015; 3(4): 823–835.