

Original Article

Assessment of Various Risk Factors Associated with Dental Implants

Vidhi Shrivastava¹, Pooran Chand², Himangi Dubey³

¹Senior Resident Faculty of Dental Sciences, ²HOD Department of Prosthodontic FODS, ³Senior Resident Department of Prosthodontic Faculty of Dental Science, KGMU Lucknow, Uttar Pradesh, India

ABSTRACT:

Background: Dental implants have been accepted as a viable treatment option for completely and partially edentulous patients. The present study was conducted to assess risk factors of associated with dental implants. **Materials & Methods:** The present study was conducted in the department of Prosthodontics. It comprised of 880 dental implants. General information such as name, age, gender etc. was recorded. Signs of peri- implantitis and fractured implant segments were considered for implant failure. **Results:** Out of 880 dental implants, 520 were in males and 360 were in females. The difference was significant (P- 0.01). Males had 45 dental implant failures and females had 40 dental implants failures. The difference was non- significant (P> 0.05). Out of 85 implant failures, 10 were seen in bruxism, 15 in autoimmune disease, 12 in diabetics, 20 in smokers and 8 in periodontal diseases. The difference was significant (P- 0.01). Implant failures were seen in type I bone (15), type II (25), type III (30) and type IV (15). The difference was non- significant (P> 0.05).

Conclusion: Dental implant failures are not uncommon. Systemic disease also associated with high implant failures. Bone quality is also risk factor for implant failures.

Key words: Bone quality, Failure, Implant.

Received: 19 January 2018

Revised: 12 February 2018

Accepted: 18 February 2018

Corresponding Author: Dr. Vidhi Shrivastava, Senior Resident Faculty of Dental Sciences, KGMU Lucknow, Uttar Pradesh, India

This article may be cited as: Shrivastava V, Chand P, Dubey H. Assessment of Various Risk Factors Associated with Dental Implants. J Adv Med Dent Sci Res 2018;6(3):61-64.

INTRODUCTION

Dental implants have been accepted as a viable treatment option for completely and partially edentulous patients. Further improvements toward the successful osseointegration of dental implants have involved modifications to both surface topography and surface chemistry. Implant design (*i.e.*, types and dimensions), surgical procedure, implant placement time, and time prior to loading have been shown to influence implant survival rates.¹

Further improvements toward the successful osseointegration of dental implants have involved modifications to both surface topography and surface chemistry. Implant design (*i.e.*, type and dimensions), surgical procedure, timing of implant placement, and time prior to loading have also been shown to influence implant survival rates.² Among patient factors, male gender, smoking, autoimmune disease, and penicillin allergy have been found to trend toward higher failure rates. In contrast,

short implants in mandibular posterior sites have been found to have a survival rate of 100%, while the same was not true for implants in the maxillary posterior position, wherein 6-mm implants had a survival rate of only 87%.

Long-term survival data, however, is required to better assess the safe and predictable use of dental implants. A few studies have reported long-term results showing more favorable survival statistics for solid screw over hollow cylinder implants, for mandibular sites over maxillary, and lower survival statistics for patients presenting with a history of periodontitis. Long-term results of implants placed with guided bone regeneration (GBR) and outcomes for the treatment of atrophic posterior maxilla have also been reported.³ The present study was conducted to assess risk factors of associated with dental implants.

MATERIALS & METHODS

The present study was conducted in the department of Prosthodontics. It comprised of 880 dental implants inserted

in patients of both genders. All patients were reviewed for 5 years and signs of implant failure were assessed. All were informed regarding the study and written consent was obtained. Ethical clearance was obtained before study.

General information such as name, age, gender etc. was recorded. Signs of peri- implantitis and fractured implant

RESULTS


segments were considered for implant failure. Results thus obtained were subjected to statistical analysis using chi-square test. P value less than 0.05 was considered significant.

Table I Distribution of implants

Males	Females	P value
520	360	0.01


Table I shows that out of 880 dental implants, 520 were in males and 360 were in females. The difference was significant (P- 0.01).

Graph I Dental implant failures




Graph I shows that males had 45 dental implant failures and females had 40 dental implants failures. The difference was non- significant (P> 0.05).

Graph II Failure rate with systemic disease



Graph II shows that out of 85 implant failures, 10 were seen in bruxism, 15 in autoimmune disease, 12 in diabetics, 20 in smokers and 8 in periodontal diseases. The difference was significant (P- 0.01).

Graph III Bone quality and failure



Graph III shows that implant failures were seen in type I bone (15), type II (25), type III (30) and type IV (15). The difference was non-significant ($P > 0.05$).

DISCUSSION

Due to recent advancements in the field of implants, there use is increasing day by day. This is now becoming the treatment of choice of missing teeth and it has got high patient compliance. However, apart from it, failures in implants are also common. It can be divided into early failure and late failure according to failure time. First, early failure is one that failed osseointegration within several weeks or several months. It was due to bone necrosis, surgical trauma, bacterial infection, inadequate initial stability and early occlusal loading. Late failure is failure that turns up after functional loading of several period of time. It takes place because of infection and excessive loading. There are many difficulties to figure out the cause of implant success and failure because it is affected by many various factors. It is hard to find a reasonable solution only with in vitro study model.⁴

In present study, out of 880 dental implants, 520 were in males and 360 were in females. We found that males had 45 dental implant failures and females had 40 dental implants failures. This is similar to Busner et al.⁵

The reasons for implants failure are lack of osseointegration during early healing, infection of the peri-implant tissues and breakage. The contraindications of implant placement are children & adolescents, epileptic patients, endocarditis, osteoradionecrosis, smoking and diabetes. Absolute contraindications consists of myocardial infarction and cerebrovascular accident, bleeding disorder, cardiac transplant, immunosuppression, active treatment of malignancy, drug abuse, and psychiatric illness, and intravenous bisphosphonate (BPs) use.⁶ In our study, out of 85 implant failures, 10 were seen in bruxism, 15 in autoimmune disease, 12 in diabetics, 20 in smokers and 8 in periodontal diseases.

David et al⁷ found that at the implant level, the cumulative survival rates at 3, 5, and 7 years were 99.3%, 99.0% and 98.4%, respectively, and at the patient level, they were 98.6%, 97.7%, and 95.9%, respectively. After adjustment to possible confounders, the multivariate analysis identified a relationship between the following risk indicators for implant failure: implant location, length and design, timing of implantation, bone grafting procedures and gender. Tissue-Level implants (n = 3863) had a very high survival rate of 99% at 3 years, which was maintained over the entire study period.

Bone-Level implants (n = 600) were as predictable with a survival rate of 99% up to 3 years, while Tapered Effect implants (n = 128) demonstrated a lower survival rate of 95% at 5 years. Short 6- mm implants in the mandibular posterior sites had a high survival rate of 100%, while in maxillary posterior positions a survival rate of only 87% was achieved. Patient factors such as smoking, autoimmune disease, and penicillin allergy were tending to associate with higher failure rates.⁸

Mittal⁹ in his study found that a total of 18 patients experienced 25 implant failures, resulting in an overall survival rate of 96.8% (2.84% and 0.38% early and late implant failures, respectively). The patient-based survival rate was 91.8%. GEE univariate and multivariate analyses revealed that a significant risk factor for implant failure was the maxillary implant. Bone grafting appeared to be a risk factor for implant failure.

CONCLUSION

Dental implant failures are not uncommon. Systemic disease also associated with high implant failures. Bone quality is also risk factor for implant failures.

REFERENCES

1. Buser D, Mericske-Stern R, Bernard JP, Behneke A, Behneke N, Hirt HP, Belser UC, Lang NP. Long-term evaluation of non-submerged ITI implants. Part 1: 8-year life table analysis

- of a prospective multi-center study with 2359 implants. Clin Oral Implants Res. 1997; 8:161-72.
2. Esposito M, Grusovin MG, Coulthard P, Thomsen P, Worthington HV. A 5-year follow-up comparative analysis of the efficacy of various osseointegrated dental implant systems: a systematic review of randomized controlled clinical trials. Int J Oral Maxillofac Implants. 2005; 20: 557-68.
 3. Albrektsson T, Branemark PI, Hansson HA, Lindstrom J. Osseointegrated titanium implants. Requirements for ensuring a long-lasting, direct bone-to-implant anchorage in man. Acta Orthop Scand. 1981; 52:155-70.
 4. Busner, Schmid J. Pathogenesis of implant failures. Periodontol. 2000; 4: 127-38.
 5. Keshmin. Patient selection and preparation. In: Branemark PI, Zarb GA & Albrektsson T (eds). Tissue integrated prosheses: Osseointegration in clinical Dentistry. Chicago: Quintessence Publishing Co. 1985; 199-209.
 6. Sharma et al. Mechanical state assessment of the implant-bone continuum: a better understanding of the Periotest method. Int J Oral Maxillofac Implants. 1995; 10: 43-9.
 7. David, Eriksson B, Lekholm U, Branemark PI, Jemt T. Longterm follow-up study of osseointegrated implants in the treatment of totally edentulous jaws. Int J Oral Maxillofac Implants. 1990; 5: 347-59.
 8. Cochran DL, Buser D, ten Bruggenkate CM, Weingart D, Taylor TM, Bernard JP, Peters F, Simpson JP. The use of reduced healing times on ITI implants with a sandblasted and acidetched (SLA) surface: early results from clinical trials on ITI SLA implants. Clin Oral Implants Res. 2002; 13: 144-53.
 9. Mittal, F. & Nisand, D. Impact of implant length and diameter on survival rates. Clinical Oral Implants Research 2006; 17: 35-51.

Source of support: Nil

Conflict of interest: None declared

This work is licensed under CC BY: *Creative Commons Attribution 3.0 License.*