

Original Article

Maxillofacial Fractures: Etiology, incidence, Pattern and Treatment of Maxillofacial Injuries in a Government Medical College of Central India

Karuna Jindwani¹, H. S. Markam², Y. K. Paharia³, Keshav Singh⁴

¹Assistant Professor, Department Of Dentistry, Shyam Shah Medical College, Rewa, ²Assistant Professor, Department Of Dentistry, Netaji Subhash Chandra Bose Medical College, Jabalpur, ³Prof. & H.O.D, Department Of Dentistry, Gajra Raja Medical College, Gwalior, ⁴Assistant Professor, Department Of Medicine, Shyam Shah Medical College, Rewa.

ABSTRACT:

Introduction: The incidence of maxillofacial fractures varies widely between various geographic locations. The large variability in reported etiology, incidence and patterns of fracture presentation are consistently influenced by geographic area, social, cultural and environmental factors. **Aims and objectives:** This descriptive study retrospectively assessed the etiology, incidence, pattern and the modalities of management of maxillofacial fractures in Netaji Subhash Chandra Bose Medical College (NSCBMC) Jabalpur, Madhya Pradesh (M. P.) for a period of one year from September 2016 till August 2017. **Materials and methods: Design:** A descriptive retrospective study. **Place and duration of study:** September 2016 to August 2017 at the department of dentistry, NSCBMC, Jabalpur (M. P.). **Methodology:** 104 consecutive patients sustaining soft and hard maxillofacial tissue injuries were included in the study. The medical records of all patients who sustained maxillofacial fractures including age, sex, etiology, influence of alcohol, pattern of facial bone fractures and the management modality were retrieved and reviewed. **Results:** As per medical records a total of 6574 patients attended the dental outpatient department (OPD). A total of 146 patients attending the dental OPD and casualty emergency room at NSCBMC, Jabalpur sustained maxillofacial injuries. Out of which, 104 patients suffered maxillofacial fractures during the period under review. Nearly 79% of the patients were men, and the most frequently affected age group was 31 to 40 years (38.46%) with males outnumbering females in all age groups. The most frequent bone fractured was mandible, accounting for 63 cases (60.57%). Road traffic accidents (RTAs) (49.03%) were the primary etiological factor followed by assault in 27.88% cases. The main stay of treatment was closed reduction with maxillo-mandibular fixation (MMF) in 58 patients. **Conclusion:** This series reflects trauma patterns at our referral centre and, as such, can provide a guide to the design of programme geared towards prevention and treatment.

Key words: Maxillofacial, Fractures, Road traffic accidents, Retrospective study.

Received: 18 February 2018

Accepted: 28 February 2018

Corresponding Author: Dr. Karuna Jindwani, Assistant Professor, Department Of Dentistry, Shyam Shah Medical College, Rewa (M.P.), India

This article may be cited as: Jindwani K, Markam HS, Paharia YK, Singh K. Maxillofacial Fractures: Etiology, incidence, Pattern and Treatment of Maxillofacial Injuries in a Government Medical College of Central India. J Adv Med Dent Scie Res 2018;6(3):101-106.

INTRODUCTION

A great diversification of human exposure to aggressive agents, make them progressive targets of numerous kinds of trauma, including maxillofacial injuries. The face, as most exposed part of the body, is particularly vulnerable to trauma.¹ and if not properly managed can negatively influence both the psychological and functional activities of the patient.² This is as a result of the centrality of the facial region as a key factor in human identity, esthetics, and general well-being.²⁻⁴

Maxillofacial trauma can be limited to superficial lacerations or abrasions of soft tissues of the face or they may involve multiple fractures of the facial bones in association with concomitant spinal and

systemic injuries thereby requiring multidisciplinary approach for their management.^{5,6} Due to cultural, environmental and socio economic factors, causes for maxillofacial trauma differ worldwide.^{7,8} Hence epidemiological studies are used to analyze the incidence and pattern of etiology, frequency, pattern and management of maxillofacial trauma.⁹ A high incidence of maxillofacial injuries due to RTAs is reported in developing nations like India while incidence due to personal violence is more in developed countries.¹⁰ Several literature reviews include the other etiological factors apart from RTA's and assault like falls, sports injuries, and animal bites and gunshot injuries. Falls

account for an important etiological factor in the facial fractures in both the very young and the elderly.¹¹

As effective injury prevention effort depends on reliable and detailed information; on the incidence, causes and pattern of injury; periodic epidemiological studies are imperative. Thus, this study was planned at the dental department of NSCB Medical College, Jabalpur to analyze the etiology, incidence and prevalence, pattern of maxillofacial fractures and for the modalities of treatment planned for the management of these fractures at our centre.

METHODOLOGY:

This comprises of a retrospective descriptive study conducted in the department of dentistry at NSCBMC, Jabalpur. It is a main referral, tertiary care centre for all places in and around the area.

Inclusion criteria:

All maxillofacial fracture patients of any age and either sex presenting to the dental department and emergency casualty from September 2016 till August 2017 were included in the study.

Exclusion criteria:

- 1) Pathology cases such as tooth ache, dento alveolar abscess, space infection, pathological fracture, temporomandibular joint dislocation other than trauma were excluded from the study.

- 2) Too old fractures with malunion.
- 3) Patients having contraindications for local and general anesthesia.

Data collection procedure:

Following due clearance from the institutional ethical review board, the case sheet records of 104 patients were thoroughly scrutinized and data regarding the study variables age, gender, etiology of trauma, diagnosis and treatment were collected on pre- designed proforma. Before enrolling the patient informed written consent was taken from subjects or their attendants. The data were interrupted using percentages wherever necessary.

RESULTS:

As per medical records a total of 6574 patients attended the dental OPD at NSCBMC, Jabalpur from September 2016 till August 2017. Out of 146 patients sustaining maxillofacial injuries, 104 patients suffered bony injuries in the form of maxillofacial fractures.

AGE AND GENDER DISTRIBUTION:

The age of the patient at the time of injury ranged from 4-55 years. Most affected age group was from 31- 40 years (38.64%) followed by patients ranging from 21 to 30 years (21.5%). The lowest frequency was observed in the age group from 51- 60 years (0.96%).

TABLE 1: AGE DISTRIBUTION OF STUDY SAMPLE (n=104)

AGE GROUP (YEARS)	NO. OF PATIENTS	PERCENTAGE
0 – 10	12	11.54
11 – 20	08	7.7
21 – 30	22	21.15
31 – 40	40	38.46
41 – 50	21	20.19
51 – 60	01	0.96
TOTAL	104	100

TABLE 2: DISTRIBUTION OF MAXILLOFACIAL ACCORDING TO GENDER (n=104)

Gender	No. of patients	Percentage
Males	82	78.84
Females	22	21.15
Total	104	100

In virtually all age- groups, more men than women were affected, the overall ratio being 3.7: 1.

ETIOLOGY OF MAXILLOFACIAL FRACTURES

TABLE 3: CAUSES OF MAXILLOFACIAL FRACTURES

CAUSES	NO. OF PATIENTS	PERCENTAGES
RTA	51	49.03
ASSAULT	29	27.88
FALL	09	8.65
SPORTS INJURY	07	6.73
MISCELLANEOUS	08	7.69
TOTAL	104	100

The most common etiology of maxillofacial injuries was RTA (51, 49.03%), followed by assault (29, 27.88%). Majority of injuries due to RTA were, as a result, of motorcycles (two- wheeler) accidents. They were either driver of vehicle or passengers and pedestrians. Alcohol by any means was found to be involved in majority of the cases. 29

(27.88%) patients fell prey to intentional injuries mainly due to assault and inter personal violence. The next common cause was due to accidental falls (09, 8.65%); sports injuries (07, 6.73%) and 08 patients (7.69%) got injured due miscellaneous causes like gunshot injuries, animal attacks, industrial accidents etc.

SITES OF FRACTURES

Fractures of the mandible were the most common type of fracture of maxillofacial skeleton (63, 60.57%). In the mandible, parasymphysis fracture was the most common type of fracture (23, 36.51%). In the midface region, most common type of fractures were fractures of zygomaticomaxillary complex (17, 41.46%).

TABLE 4: SITE OF MAXILLOFACIAL FRACTURES

SITE	NO. OF CASES	PERCENTAGES
MANDIBLE	63	60.57
MIDFACE	41	39.42
A) LEFORT B) ZYGOMATICOMAXILLARY C) NASO- ETHMOID		
TOTAL	104	100

TABLE 5: ANATOMICAL LOCATIONS OF MANDIBULAR FRACTURE

LOCATION	NO. OF CASES	PERCENTAGES
DENTOALVEOLAR	03	4.76
SYMPHYSIS	12	19.04
PARASYMPHYSIS	23	36.51
BODY	08	12.69
ANGLE	05	7.93
CONDYLE	07	11.11
RAMUS	03	4.76
CORONOID	02	3.17
TOTAL	63	100

TABLE 6: ANATOMICAL LOCATION OF MIDFACE FRACTURES

LOCATIONS	NO. OF CASES	PERCENTAGES
DENTOALVEOLAR	05	12.19
LEFORT I	03	7.31
LEFORT II	09	21.95
LEFORT III	04	9.75
ZYGOMATICOMAXILLARY	17	41.46
NASOETHMOID	03	7.31
TOTAL	41	100

TREATMENT PROTOCOL:

Several methods of reduction and fixation were used in the treatment of mandible fractures. Of the total 104 fractures, 58 cases (55.77%) were treated by closed reduction with arch bars and MMF. Splint fixation was used in 2 cases (1.92%) in a mandibular body fracture of a 4 years old child and in a double fracture of mandible in a 55 years old edentulous denture wearing patient. Most of the patients (34 cases, 32.69%) with mid face fractures and pan facial trauma cases were treated with open reduction and internal fixation (ORIF) with bone miniplates under general anesthesia (G. A.). All such cases were supported by MMF. However, in internal rigid fixation, early removal of MMF was advised. The plain Ehrich's arch bar was used to manage dentoalveolar fractures involving maxillary and mandibular teeth in 8 patients (7.69%).

TABLE 7: DISTRIBUTION OF TREATMENT MODALITIES

TREATMENT	NO. OF CASES	PERCENTAGES
ARCH BAR FIXATION	08	7.69
SPLINTS AND WIRING	02	1.92
CLOSED REDUCTION (MMF)	58	55.77
ORIF (BONE PLATES + MMF)	34	32.69
CONSERVATIVE	02	1.92
TOTAL	104	100

2 patients (1.92%) with trismus due solitary fracture of coronoid process of mandible were managed by conservative therapy of restrictions in diet and regular follow up for monitoring of increase in mouth opening and balancing of occlusion bilaterally.

DISCUSSION:

Fractures of the facial skeleton are a common finding following trauma and therefore form an integral part of the overall duty of an oral and maxillofacial surgeon.¹² The result of epidemiological surveys on the causes and incidence of maxillofacial fractures tend to vary with geographic region, socioeconomic status, culture, religion and era.^{6,11,13,14}

Site and age distribution:

The maxillofacial fractures occur most commonly in young adults especially males in the third and fourth decades of life because they are majorly involved in many outdoor activities, take part in dangerous exercises and sports, and are most likely to be involved in violence.^{5,12,15} In our study 78.84% patients were males with the male: female ratio of 3.7: 1. The sex ratio in various studies range from 2.3: 1¹⁶ to 11.8: 1.¹⁷ Injured males predominantly in their fourth decade (31- 40 years) reported to our centre. This differs slightly from most of the studies which considered injuries commonly reported in the third decade of life.^{6,16,18,19} And contrasts with the report of Karyouti,¹⁷ who gave the age group of 0- 5 years as having the highest incidence. This lowest frequency was observed in the age group above 50 years (0.96%) which was most probably due to limited outdoor activities in old age.

Etiology of trauma:

In coincidence with the changes in the community lifestyle, industrialization, transportation and legislative measures, the causes of maxillofacial fractures also tend to change. In most previous epidemiological studies accidents were the most common cause of maxillofacial fractures,^{8,14,17,20-22} and the present study supports these findings with 49.03% of the patients were injured in RTA's. However, in contrast to other studies carried out in developed countries, which reported assaults as the most common cause of maxillofacial injuries.^{5,8,12,13,19} The reasons for this high rate of RTA in India includes poor road networks, improper licensing of drivers/ riders, no usage of seat belts, over speeding, overloading underage driving, neglect of helmets by motorbike riders and non compliance with traffic rules.^{5,12} Alcohol and unemployment are also contributing factors which has commonly featured in other similar studies.²³⁻²⁵ Other etiological factors reported in this study include falls from height and sports- related injuries, gunshot injuries, animal attacks and industrial accidents.

Sites and patterns of fractures:

When the maxillofacial region is injured, the mandible is more vulnerable than the mid face fractures.¹⁴ This could be because the mandible is mobile and has less bony support than the well articulated midfacial bones. Mandibular fractures were the highest occurring fractures in the present review accounting for 60.57% of the total fractures. This is in agreement with other reports^{8, 11, and 12} but differs from studies from the western world where nasal bone fractures⁵ and zygomatic complex fractures²⁶

occur more frequently. In this study para- symphysis (36.51%) was the most common site involved in the injury similar to few other studies^{8, 11 and 27} and differed from other some other studies which reported more frequent mandibular body fractures^{5, 12} and condylar neck fractures.²⁸ In the middle third, the zygomatico maxillary complex is the most involved site due to its projection and multiple articulations with other facial skeletons making it very vulnerable to fractures on impact.^{13,18,26}

Treatment protocol:

The age old principle of fracture management; reduction and immobilization also applies to maxillofacial region; however, the pathway to achieving this principle is influenced by many factors.⁴ It should be noted that the treatment outcome of maxillofacial fractures is mainly dependent among other things on the degree of injury, type of fracture, the expertise of the surgeon, and available technology.²⁻⁴ In the past 20 years, changes in maxillofacial trauma management have been strongly influenced by innovations in materials and technology,²⁹ since objectives such as early recovery, segment stability and patient comfort have been considered paramount in the treatment of maxillofacial fractures.³⁰ Thus, the management of maxillofacial injuries is a real challenge for any oral and maxillofacial surgeon and demands both skill and expertise. In our study closed reduction including MMF with the help of Ehrlich's arch- bar was done in 58 fractures (55.77%) which coincides with other reports from literature where closed reduction was the most frequently used treatment modality for facial injuries.^{4,28,29} All panfacial fractures (34 cases, 32.69%) multiple fractures with occlusal derangement and displaced fractures that were not reduced by closed reduction and unstable ZMC fractures after elevation were treated by open reduction and direct fixation by miniplate osteosynthesis. The place of ORIF with miniplate osteosynthesis in the surgical management promises a shortened period of MMF, bony union with minimal callus formation, rapid recovery of normal jaw functions, and maintenance of normal body weights among others.^{4,31} Despite these advantages, ORIF of facial fractures has not become popular at our centre like in most developing countries mainly due to its cost and the time required to procure the plates.^{4,32} And moreover simple methods of fixation give satisfactory results. However, advantages and effectiveness versus the cost of miniplates fixation need further survey in our region.

LIMITATIONS:

The major limitation of our survey is its retrospective design, mainly due to poor record keeping and insufficient data from the available patient's records. However, we tried to present an analysis based on prospective- database, although retrospectively analysed and it is for the first time that epidemiology and pattern of facial injuries have been brought into limelight in our environment. To overcome the limitation, a larger- data based prospective study may help to improve upon our

present findings and also discover subsequent changes in the trends and pattern of these fractures.⁴

CONCLUSION AND RECOMMENDATIONS:

Present study supports that regular epidemiologic evaluations of maxillofacial fractures allow a detailed analysis of facial injuries in our environment, providing important support to install clinical and research priorities, since risk factors and patterns of can be identified.³³ From our study, it seems reasonable to assume that RTA's remain the leading cause of maxillofacial fractures and is closely followed by assault especially among men in their productive years. Mandibular fracture was identified as the most frequently occurring bony injury on face with specific preponderance to the parasymphysis region in 36.5% of all lower jaw fractures. Even though miniplate osteosynthesis is now the 'gold standard'³³ of fracture management across the surgical world, a high proportion of our patients were treated by traditional methods of closed reduction and MMF.

In conclusion, this study provides relevant data on patterns of maxillofacial and dental trauma in a specific population. In light of this study, following recommendations are advised to reduce the maxillofacial trauma in this belt and improvise the treatment modalities-

- 1) Amendments in the traffic rules and legislations about seat belt and helmet usage followed by their strict implementation on roads.
- 2) Regular and timely maintenance of faulty roads by government.
- 3) Strict implementation of legislations against over loading in public vehicles and over speeding in youth.
- 4) Ensuring disposal of out of date vehicle by government.
- 5) Provision of pedestrian friendly paths and segregation of heavy and light motor vehicles.
- 6) Legal prohibition of drunk and driving and usage of cell phones while driving.
- 7) Improvement in education and socio economic set up aimed to reduce the incidence of fractures secondary to assaults.
- 8) Adequate safety features in building design in order to reduce the incidence of fractures secondary to fall in children.
- 9) Educational campaigns for recommending the use of mouth guards, helmets, knee and elbow-pads while practicing sports.

Thus, our efforts via this survey emphasize the need of our government to enforce the mentioned preventive measures in order to bring a steady decrease in facial injuries in our environment. The government is also hereby welcomed to enforce the health insurance scheme and ensure a good and secure environment for the marketing, distribution and prompt availability of bone plates at a tertiary care centre like ours in a developing country.

REFERENCES:

1. Oburanya AA, Adeleye AO, Aladelusi TO, Fasola AO. Updates on the epidemiology and pattern of traumatic maxillofacial injuries in a Nigerian university teaching hospital: a 12- month prospective cohort in- hospital outcome study. *Craniomaxillofac Trauma Reconstr.* 2015; 8(1): 50- 58, doi: 10.1055/ 5-0034-1384740.
2. K Yamamoto, Y. Matsusue, S. Horita, K. Murakami, T. Suigiura, and T. Kirita, "Clinical analysis of midfacial fractures," *Materia Socio Medica*, vol.26, no.1, pp, 21- 25, 2014.
3. A. Parashar and R. K. Sharma, "Unfavorable outcomes in maxillofacial injuries: how to avoid and manage," *Indian Journal of Plastic Surgery*, vol. 46, no. 2, pp. 221- 234, 2013.
4. Udeabor S. E, Akinbani. B. O., Yarhere. K. S, Obiechina AE, Maxillofacial fractures: Etiology, pattern of presentation, and Treatment in university of Port Harcourt teaching hospital, Port Harcourt, Nigeria. *Journal of Dental Surgery.* 2014; dx.doi.org/ 10.1155/ 2014/ 850814.
5. E. D. Arshan, A. G. Solakoglu, E. Komut et al., "Assessment of maxillofacial trauma in emergency department," *World Journal of Emergency Surgery*, vol9, no. 1, article 13, 2014.
6. Khan M. Management of maxillofacial trauma. *AFID Dent J* 1998; 10: 18- 21.
7. Mock C, Quansab R, Krishnan R, Arreola- Rise C, Rivara F. Strengthening the prevention and care of injuries worldwide. *Lancet*, 2004; 363: 2172- 2179.
8. Fasola AO, Nyako EA, Obiechina AE, Arotiba JT. Trends in the characteristics of maxillofacial fractures in Nigeria. *J Oral Maxillofac Surg* 2000; 61: 1140- 43.
9. Gassner R, Tuli T, Hachi O, Rudish A, Ulmer H. Cranio- Maxillofacial trauma: a 0 year review of 9, 543 cases with 21,067 injuries. *J Craniomaxillofac Surg* 2003; 31: 51- 61.
10. Kyrgides A, Koloutsos G, Kommata A, Lazarides N, Antoniadis K. Incidence, aetiology, treatment out come and complications of maxillofacial fractures. A RETROSPECTIVE STUDY FROM NORTHERN GREECE. *Craniomaxillofac Surgery.* 2013; 41(7): 637- 643. Doi: 10.1016/ j.jcms. 2012. 11. 046.
11. Banks P. Killey's fractures of the middle third of the facial skeleton 5th ed. Oxford: Butterworth Heinemann 1988.
12. W. L. Adeyeno, A. L. Ladeinde, M. O. Ogunlewe, and O. James," Trends and characteristics of oral and maxillofacial injuries in Nigeria: a review of the literature", *Head & Face Medicine*, Vol 1 article 7, 2005.
13. Schiender D, W Keammerer, Scheon G, Dinu C, Radloff S Bchorer R. Etiology and injury patterns of maxillofacial fractures from the years 2010 to 2013 in Mecklenburg- Western Pomerania. Germany: A retrospective study of 409 patients. *J Cranio Max Fac Surgery* 2015; 43: 1948- 51.
14. Boffano P, Rocciia F, Zavaltero E, Dediol E, Uglesic V, Kovacic Z, et al. European Maxillofacial trauma (EURMAT) project. A multicentre and prospective study. *J Craniomaxillofac Surg.* 2015; 43(1): 62- 70. Doi: 10.1016/ j.jcms.2014. 10. 011.
15. B Van den Bergh, K. H. Karagozoglu, M. W. Heymans, and T. Forouzanfar, "Aetiology and incidence of maxillofacial trauma in Amsterdam: a retrospective analysis of 579 patients." *Journal of Cranio- Maxillofacial Surgery*, Vol. 40, no. 6, pp. e165- e169, 2012.
16. Banjec Y, Lownie J F, Cleaton Jones PE, Lownie MA, Maxillofacial injuries in a group of South Africans under 18 years of age. *Br. J Oral Maxillofac Surg* 1996; 34: 298- 302.

17. Iida S, Kogo M, Suguiura T, Mima T, Matsya T. Retrospective analysis of 1502 patients with facial fractures. *Int J Oral Maxillofac Surg* 2001; 30: 286- 90.
18. Mesgarzadeh AH, Shahamfar M, Azar SF, Shahanfar J . Analysis of the pattern of maxillofacial fractures in Noerth Western of Iran. A retrospective study. *J Emerg Trauma Shock*. 2011; 448- 52.
19. Kapoor P, Kalra N. A. A retrospective analysis of maxillofacial injuries reporting to a tertiary care hospital in East Delhi. *Int J Crit Inj Sci* 2012; 206- 10.
20. Moore BK, Smit R, Colquhoun A, Thompson WM. Maxillofacial fractures at Waikato Hospital, New Zealand: 2004 to 2013. *NZ Med J* 2015; 128(1426) 96- 102.
21. Venugopal MG, Sinha R, Menon PS, Chattopadhyay PK, Roy Chowdhury SK. Fractures in the maxillofacial region: A four- year retrospective study. *MUFAI* 2010; 66: 14- 17.
22. Chandra- Shekhar BR, Ready CV. A five year retrospective statistical analysis of maxillofacial injuries in patient admitted and treated at two hospitals of Mysore city. *Indian J Dent Res*. 2008; 19: 304- 12.
23. Chrcanovic BR, Abreu MH, Freire- Maid B, Souza LN. Facial fractures in children and adolescents: a retrospective study of 3 years in a hospital in Belo Horizonte, Brazil. *Dent Tramadol*, 2010 Jun; 26 (3): 262- 70. Epub 2010 Apr 23.
24. Alvi , Doherty T, Lewen G: Facial fractures and concomitant injuries in trauma patients. *Laryngoscope* 2003, 113: 102- 6.
25. O' Meara C, Witherspoon R, Harpangama N, Hyam DM, Alcohol and interpersonal violence may increase the severity of facial fractures. *Br J Oral Maxillofac Surg* 2012; 50: 36- 40.
26. A. Dibaie, S. Raissian, and S. Ghafarzadeh, " Evaluation of maxillofacial traumatic injuries of forensic medical centre of ahwaz, Iran, in 2005", *Pakistan Journal of Medical Sciences*, Vol 25, no. 1, pp. 79- 82, 2009.
27. Kotecha S, Scannell J, Monaghan A, William RW. A four year retrospective study of 1062 patients presenting with maxillofacial emergencies at a specialist paediatric hospital. *Br J Oral Maxillofac Surg*. 2008; 46: 293- 6.
28. Ahmed HEA, Jaber MA, Fanas SHA, Karas M. The pattern of maxillofacial fractures in Sharjah, United Arab Emirates; a review of 230 cases. *Oral Surg Oral Med Oral Pathol Oral Radio Endod* 2004; 98: 166- 70.
29. Adebayo ET, Ajike OS, Adekye EO. Analysis of the pattern of maxillofacial fractures in Kaduma, Nigeria. *Br J Oral Maxillofac Surg* 2003; 41: 396- 400.
30. Gabrielli MAC, Gabrielli MFR, Marcantonio E, Holuchi – Vieira E. Fixation of mandibular fractures with 2.0 mm miniplates review of 191 cases. *J Oral Maxillofac Surg* 2003; 61: 430- 36.
31. I. Abbas and K. Ali, " Management of mandibular fractures- a prospective study, " *Pakistan Oral & Dental Journal*, vol.22, no. 2, pp. 151-152, 2002.
32. Ahmed S, Punjabi SK, Rehman HV, Aslam MA, Haider SM. Analysis of maxillofacial injuries spread over one year period in Karachi sample. *Pak Oral Dent J*. 2014; 34(1): 50- 53.
33. Jindwani K, Paharia. Y. K, " Demographic distribution, etiological spectrum, injury characteristics, management and preventive recommendations for maxillofacial injuries at a tertiary care centre in central India. *Journ of Evol of Med and Dent Sci* 2015; vol. 4, issue 64, Aug 10; pp 11193- 11201, DOI 10. 14260/ jemds/ 2015/ 1612.

Source of support: Nil

Conflict of interest: None declared

This work is licensed under CC BY: *Creative Commons Attribution 3.0 License*.