

Original Article

Assessment of Platelet count changes in patients with Chronic Periodontal Diseases

Anju Gautam¹, Awadhesh Chandra Nagar², Neeraj Kumar Dhiman³

¹Assistant Professor, Department of Periodontics, ³Assistant professor, Oral and Maxillofacial surgery, Faculty of Dental sciences, Institute of Medical Sciences, Banaras Hindu University, Varanasi.

²Medical Officer, Sir Sunderlal Hospital, Institute of Medical Sciences, Banaras Hindu University, Varanasi, U.P.

ABSTRACT

Objectives: The objective of the study was to determine whether platelets count were changed in patients with chronic periodontitis as compared to healthy subjects. **Methodology:** A total of 154 systemically healthy subjects (73 periodontally healthy and 81 with chronic periodontitis) of age group 25-50 years were selected. Periodontal parameters including Plaque Index (PI), Gingival Index (GI), Pocket Depth (PD), Clinical Attachment Level (CAL) were measured. Venous blood samples were taken to analyze the platelet count. **Result and Conclusions:** The platelet count was higher in subjects with chronic periodontitis as compared to healthy but the result was statistically non significant.

Key words: Periodontitis, platelets count, hematological parameter.

Received: 10 October 2018

Revised: 22 October 2018

Accepted: 25 October 2018

Corresponding Author: Dr. Anju Gautam MDS, Assistant Professor, Department of Periodontics, Faculty of Dental sciences, Institute of Medical Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh, India

This article may be cited as: Gautam A, Nagar AC, Dhiman NK. Assessment of Platelet count changes in patients with Chronic Periodontal Diseases. J Adv Med Dent Scie Res 2018;6(11):27-29.

INTRODUCTION

Periodontal disease is an inflammatory infectious diseases caused by the interaction of plaque bacteria and the host leading to destruction of the alveolar bone and connective tissue supporting the tooth. The primary etiology of periodontal diseases is a group of specific periodontal pathogens, and the host's immunologic and inflammatory response to the bacteria that result in periodontal destruction (Listgarten 1986).¹

Systemic factors modify the periodontal diseases mainly by acting on the normal immune and inflammatory defense cells.

In the past, studies have been conducted regarding whether periodontitis may affect the changes in cellular and molecular components of blood circulation (Loos 2005).²

The studies suggest that the patients with periodontal diseases have elevated levels of systemic inflammation markers including leukocytes (Danesh et al.,1998, Loos et al., 2000)^{3,4} neutrophils, and and C-reactive protein (Schreiber et al.,1982, Loimaala et al.,2006)^{5,6} as compared to healthy individuals. Papapanagiotou et al. have shown that platelets activation have increased in

patients with periodontitis compared with platelets from healthy controls.⁷

The current study was conducted to assess the platelet count in blood of systemically healthy individuals and chronic periodontitis patients.

MATERIALS AND METHODS

The subjects for the study were selected from patients reporting to the Outpatient Department of Periodontics, Faculty of Dental Sciences, Banaras Hindu University (BHU), Varanasi. The study comprised of 154 subjects, inclusive of both sexes, aged between 25 and 50 years, selected on a random basis. The purpose of the study and methodology were explained to the subjects. Those participants who gave written informed consent for the study were included in the study.

Patients with any systemic diseases, pregnant women, those under medication within the past 6 months, smokers and all subjects who have received any periodontal treatment within 6 months before examination, and those having conditions which could aggravate periodontal manifestations were excluded from the study.

Periodontal examination was carried out after selection of subjects. Subjects were divided into two groups based on

the periodontal examination: group I (healthy participant), and Group II (patients with periodontitis). Periodontal examination consisted of Plaque Index (PI), Gingival Index (GI), Pocket Depth (PD), Clinical Attachment Level (CAL). Healthy subjects displayed periodontal pocket depth <3 mm, Clinical attachment loss <3 mm, with no clinical sign of gingivitis. Periodontitis patients exhibited ≥ 4 mm of pocket depth and Clinical attachment loss ≥3 mm in more than 30% sites assessed in the oral cavity.

2 ml of venous blood sample were obtained by venipuncture method in the antecubital fossa and stored in vials containing EDTA. The platelet counts were determined by using an automated cell counter within 24 hours after sample collection, at the hematology laboratory of the institution.

Data were analyzed using statistical software, namely, Statistical Package for the Social Sciences (SPSS, Version 16, IBM Analytics). Student t-test was used to analyse the differences between the two groups and P < 0.05 was considered statistically significant.


RESULTS

Table 1 shows the mean values of platelet counts in subjects of both the groups. The mean platelet count in the subjects with chronic periodontitis (253.89±52.87 × 10⁹ cells/L) was higher as compared to the healthy group (239.07±56.93 × 10⁹ cells/L). The mean platelet counts are graphically represented in fig 1. The observed difference between the two groups was found to be statistically non-significant.

Table 1: Comparison of Platelet count between groups.

Group	N	Platelet count (x10 ⁹ /L) Mean±SD	P Value
Group I (Healthy)	73	239.07±56.93	0.0960
Group II (Periodontitis)	81	253.89±52.87	

Figure 1: Bar diagram showing Platelet count in periodontitis and healthy subjects



DISCUSSION

Chronic periodontitis is an infectious disease of the supporting structures of the teeth and is caused by complex, anaerobic, gram-negative specific periodontal bacteria. Periodontal destruction may results from the action of toxic products released from the specific pathogenic bacteria, as well as from the host responses initiated against the periodontal bacteria and their products. The studies suggest that the periodontal diseases increase the risk of systemic conditions like cardiovascular diseases, cerebrovascular diseases, atherosclerosis, and diabetes mellitus.

A total hematological analysis is frequently used to assess the presence of systemic infection or inflammation, and the question arises whether periodontal infections can also affect the hematological parameters such as the differential counts of white blood cells, red blood cells, and platelets count. Some of the studies have demonstrated differences in the hematological parameters in subjects with chronic periodontitis as compared to healthy subjects (Lopez et al., 2012).⁸

In the present study, the mean platelet count in the subjects with chronic periodontitis was 253.89±52.87 × 10⁹ cells/L and in healthy subjects was 239.07±56.93 × 10⁹ cells/L. The mean platelet count in the subjects with chronic periodontitis was higher as compared to the healthy group but the difference between the two groups was found to be statistically non-significant. The result of this study is in agreement with the previous study done by Al-Rasheed A. and Nicu et al.^{9, 10} This is similar to the previous studies showing an increase in platelet levels and activation in periodontitis patients group (Papapanagioutou et al., 2009⁷; Wakai et al., 1999)¹¹ and the study that demonstrated periodontal treatment results in a lowering of platelet count (Christan et al., 2002).¹² In a study conducted by Lourbakos et al,¹³ It has been concluded that dental plaque bacteria, including the periodontal pathogen Porphyromonas gingivalis, induces platelet activation and aggregation.

CONCLUSIONS:

From the results of the present study, it can be concluded that chronic periodontitis may elevate the levels of platelet counts as compared to healthy control patients and cab be considered as one of the mechanisms which explain that periodontitis is linked to the systemic inflammatory conditions.

REFERENCES:

- Listgarten MA. A perspective on periodontal diagnosis. J Clin Periodontol 1986;13:175-81.
- Loos BG. Systemic markers of inflammation in Periodontitis. J Periodontol 2005;76: 2106-2115.
- Danesh J, Collins R, Appleby P, Peto R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: Meta-analyses of prospective studies. JAMA 1998; 279: 1477-1482.
- Loos BG, Craandijk J, Hoek FJ, Wertheim-van Dillen P, M.van der Velden U. Elevation of systemic markers related to cardiovascular diseases in the peripheral blood of periodontitis patients. J Periodontol 2000;71:1528-1534.

5. Schreiber G, Howlett G, Nagashima M, Millership A, Martin H, Urban J, Kotler L. The acute phase response of plasma protein synthesis during experimental inflammation. *J Biol Chem* 1982;257:10271–10277.
6. Loimaala A, Rontu R, Vuori I, Mercuri M, Lehtimäki T, Nenonen A, Bond MG. Blood leukocyte count is a risk factor for intima media thickening and subclinical carotid atherosclerosis in middle-aged men. *Atherosclerosis* 2006;188:363–369.
7. Papapanagiotou D, Nicu EA, Bizzarro S, Gerdes VE, Meijers JC, Nieuwland R, van der Velden U, Loos BG. Periodontitis is associated with platelet activation. *Atherosclerosis* 2009;202: 605–611.
8. López R, Loos BG, Baelum V. Hematological features in adolescents with Periodontitis. *Clin Oral Invest* 2012;16:1209–1216.
9. Abdulaziz Al-Rasheed. Elevation of white blood cells and platelet counts in patients having chronic periodontitis. *Saudi Dent J*. 2012 Jan; 24(1): 17–21.
10. Nicu EA, Velden UVD, Nieuwland R, Everts V, Lo O S BG. Elevated platelet and leukocyte response to oral bacteria in periodontitis. *Journal of Thrombosis and Haemostasis*, 7: 162–170.
11. Wakai K., Kawamura T., Umemura O., Hara Y., Machida J., Anno T., Ichihara Y., Mizuno Y., Tamakoshi A., Lin Y., Nakayama T., Ohno Y. Associations of medical status and physical fitness with periodontal disease. *J. Clin. Periodontol.* 1999;26(10):664–672.
12. Christan C., Dietrich T., Hägewald S., Kage A., Bernimoulin J. White blood cell count in generalized aggressive periodontitis after non-surgical therapy. *J. Clin. Periodontol.* 2002;29(3):201–206.
13. Loubakos A., Yuan Y., Jenkins A., Travis J., Andrade-Gordon P., Santulli R., Potempa J., Pike R.N. Activation of protease-activated receptors by gingipains from *Porphyromonas gingivalis* leads to platelet aggregation: a new trait in microbial pathogenicity. *Blood*. 2001;97(12):3790–3797.

Source of support: Nil

Conflict of interest: None declared

This work is licensed under CC BY: ***Creative Commons Attribution 3.0 License.***