

Original Research

To study the clinical profile of anaemia in a tertiary care hospital

Jagda Nand Jha

Asst Prof, Department of General Medicine, Noida International Institute of Medical Sciences, Greater Noida, India

ABSTRACT:

Aims: To study the clinical profile of anaemia in a tertiary care hospital. **Methods:** This was a prospective observational study conducted in the Department of General Medicine. In all 100 patients thorough history was taken, general physical examination and systemic examination were done. Patients were subjected to routine blood investigations including complete blood count, peripheral smear study and serology for viral markers. **Results:** 48(48%) were easy fatigability and generalised weakness the most common symptoms of anaemia in our study. Incidentally detected patients constituted 31% of patients and were the second most common in occurrence. This was followed by breathlessness seen in 19% of patients. Pallor was noted in all patients. Platonychia/koilonychia suggesting iron deficiency anaemia was seen in 26(26%) of patients, whereas knuckle pigmentation suggestive of megaloblastic anaemia was observed in 18(18%) of patients. 15 (15%) patients presented with anaemia in failure as evidenced by elevated jugular venous pulse and pedal oedema. On systemic examination haemic murmurs on CVS examination were detected among 22(22%) patients. Severe anaemia showed 84 (84%) highest occurrence. Microcytic hypochromic anaemia 50(50%) attributed to iron deficiency unless proved otherwise was the most common form of anaemia in our study. Dimorphic anaemia 27(27%) was the second most common suggesting that nutritional anaemia continues to predominate in our part of world. **Conclusion:** Nutritional anaemia particularly iron deficiency anaemia is the most common cause of anaemia. It tends to affect the working age group and females predominantly. Patients continue to present with severe anaemia to the hospital.

Keywords: Anaemia, Clinical profile, Laboratory profile

Received: 17 July, 2021

Accepted: 27 August, 2021

Corresponding author: Jagda Nand Jha, Asst Prof, Department of General Medicine, Noida International Institute of Medical Sciences, Greater Noida, India

This article may be cited as: Jha JN. To study the clinical profile of anaemia in a tertiary care hospital. J Adv Med Dent Scie Res 2021;9(9):77-80.

INTRODUCTION

Anemia is considered the most prevalent nutritional deficiency globally, affecting about a quarter of the world population especially children and women of reproductive age.¹ Anemia is generally defined as a reduction in red cell mass or blood hemoglobin concentration characterized by decreased oxygen carrying capacity of blood which results in tissue anoxia producing various signs and symptoms. Anemia is not a diagnosis in itself but merely an objective sign of presence of disease. Anemia in children differs from those of adults as they tend to be more pronounced and develop rapidly. As much as 51% children in 0-4 years and 46% children 5- 12 years are anemic in developing regions.^{2,3}

Anaemia is an indicator of poor nutrition and poor health. It is a global public health problem affecting both developed and developing nations. In 2002 iron deficiency anaemia was considered amongst the most

important contributing factor to the global burden of disease.⁴ India is among the countries with highest prevalence of Anaemia in the world. It is estimated that about 20%-40% of maternal deaths in India are due to Anaemia; India contributes to about 50% of global maternal deaths due to Anaemia.⁵ Anaemia are of different types. Iron deficient Anaemia is the most common type of anaemia. Quite frequently faulty nutrition is the cause of Anaemia. There are many factors like inadequate diet, unsatisfactory method of preparation of food, faulty social habits, unhygienic practices, associated infections and infestations contributing to the causation of nutritional Anaemia.^{6,7} India lies partly in the tropics and partly in subtropics with extreme variations of climate. In the region where hot and humid climate prevail throughout the best part of the year, the loss of iron through sweat is appreciable. Iron is lost through sweat to the extent of 15mg per month. This suggests dermal loss of iron

should be one of the possible contributing factors in the genesis of iron deficiency Anaemia in the tropics.^{6,7} Our study aimed to provide clinical and laboratory profile of anaemia patients at a tertiary care hospital.

MATERIAL AND METHODS

This was a prospective observational study conducted in the Department of General Medicine, after taking the approval of the protocol review committee and institutional ethics committee. 100 patients with more than or equal to 18 years of age of both sexes and Patients with anaemia as per WHO definition were

included in this study. Patients not willing to give informed consent were excluded from the study.

In all 100 patients thorough history was taken, general physical examination and systemic examination were done. Patients were subjected to routine blood investigations including complete blood count, peripheral smear study and serology for viral markers. Required radiological investigations were done and further studies like bone marrow examination, iron profile, vitamin B12 and folate levels were done in selected patients who did not respond to therapy started based on peripheral smear report. Stool for occult blood was done among elderly patient presenting with iron deficiency anaemia

RESULTS

Among 100 patients studied 30(30%) were males and 70(70%) were females. In our study it was found that anaemia had its highest occurrence in the age group of 20-30 years 55(55%) followed by below 20 year age group 22(22%). It was least among individuals aged above 50 years 1(1%) Table 1.

Table 1. Demographic profile of the patients

Gender	N=100(%)
Female	70(70%)
Male	30(30%)
Age (years)	
Below 20 years	22(22%)
20-30years	55(55%)
30-40years	13(13%)
40-50years	9(9%)
Above 50years	1(1%)

48(48%) were easy fatigability and generalised weakness the most common symptoms of anaemia in our study. Incidentally detected patients constituted 31% of patients and were the second most common in occurrence. This was followed by breathlessness seen in 19% of patients (Table 2).

Table 2: Symptoms of anaemia patients

Symptoms	Number of patients
Easy fatigability and generalised weakness	48 (48%)
Breathlessness	19 (19%)
Swelling of limbs, puffiness of face	7 (7%)
Giddiness	10(10%)
Chest pain	2 (2%)
Fever	14 (14%)
Tinnitus	5 (5%)
Asymptomatic (incidentally detected)	31 (31%)

Pallor was noted in all patients. Platonychia/koilonychia suggesting iron deficiency anaemia was seen in 26(26%) of patients, whereas knuckle pigmentation suggestive of megaloblastic anaemia was observed in 18(18%) of patients. (Table 3).

15 (15%) patients presented with anaemia in failure as evidenced by elevated jugular venous pulse and pedal oedema. None of the patients in this study was due to hemolysis. Hence icterus seen in 7(7%) patients was due to ineffective erythropoiesis seen in patients with megaloblastic anaemia. On systemic examination haemic murmurs on CVS examination were detected among 22(22%) patients. Bibasilar crepts not attributable to other diseases were found among 5 patients. Isolated hepatomegaly was found in 12(12%), splenomegaly in 7(7%) and hepatosplenomegaly was found in 7 patients.

Table 3: Signs in patients with anaemia

Signs	Number of patients
Tachycardia	49 (49%)
Tachypnea	17(17%)
Elevated JVP	15 (15%)
Pallor	100 (100%)
Icterus	7 (7%)

Pedal oedema	13(13%)
Platonychia/koilonychia	26 (26%)
Knuckle pigmentation	18(18%)

Table 4: Degree of anaemia

Degree	Number of patients
Mild anaemia	0 (0%)
Moderate anaemia	16(16%)
Severe anaemia	84(84%)

On laboratory examination degree of anaemia (as defined by WHO) was distributed as shown in Table 4. None of the patients admitted in the hospital had mild anaemia (defined as Hb between 11-11.9 g/dl in women and 11-12.9 g/dl in men aged 15 years or more). Moderate anaemia (defined as Hb between 8 to 10.9 g/dl in both males and females) was seen in 16% of patients. Whereas severe anaemia (defined as Hb less than 8 g/dl in both males and females) showed 84 (84%) highest occurrence (Table 4). Microcytic hypochromic anaemia 50(50%) attributed to iron deficiency unless proved otherwise was the most common form of anaemia in our study. Dimorphic anaemia 27(27%) was the second most common suggesting that nutritional anaemia continues to predominate in our part of world (Table 5).

Table 5: Peripheral smear study in patients with anaemia

Peripheral smear	Number of patients
Microcytic hypochromic anaemia	50(50%)
Macrocytic anaemia	2 (2%)
Dimorphic anaemia	27 (27%)
Normocytic normochromic anaemia	21 (21%)

DISCUSSION

In our study it was found that anaemia had its highest occurrence in the age group of 20-30 years 55(55%) followed by below 20 year age group 22(22%). It was least among individuals aged above 50 years 1(1%). predominantly affecting the working class of the population. Similar observations were made in a study conducted by Azad KL et al.⁸ Statistically 70% of patients were females and rest were males depicting a female preponderance. Such female dominance was also shown in studies conducted by Alvarez-Uria G et al, and Talwelkar SR et al.^{9,10} WHO statistics noted that the prevalence of iron deficiency anaemia, most common cause of anaemia in females in the age group of 15-49 years is 52%.¹¹ This study upholds this fact as well. In our study 48 (48%) were easy fatigability and generalised weakness the most common symptoms of anaemia in our study. Incidentally detected patients constituted 31% of patients and were the second most common in occurrence. Easy fatigability as the predominant symptom was also noted in studies conducted by Dashratham P et al, and Gayathri BN et al.^{12,13} Incidentally detected anaemia constituted the second most common class. This may be explained by the fact of lack of knowledge or presence of chronic anaemia. 19 (19%) presented with breathlessness whereas puffiness of face and swelling of limbs was seen in 7 (7%) of patients. Fever secondary to anaemia and not attributable to any other cause was seen in 14(14%) patients. Fever as a symptom of anaemia was also noted in study conducted by S. Selvamuthukumar.¹⁴ Anaemia causing tinnitus after ruling out neuro-otologic and other secondary causes was seen in 5(5%) patients.

As far as signs on general physical examinations were concerned pallor was the universal finding present in 100% of patients. Such predominance of pallor as a sign on examination was noted in studies conducted by Gayathri BN et al, and Vineetha et al.^{13,15} This was followed by tachycardia seen in 49(49%) patients. 15(15%) patients presented with anaemia in failure as evidenced by elevated jugular venous pulse and pedal oedema. None of the patients in this study was due to hemolysis. Hence icterus seen in 7(7%) patients was due to ineffective erythropoiesis seen in patients with megaloblastic anaemia. Signs depicting the etiology i.e platonychia/koilonychia suggesting iron deficiency anaemia and knuckle pigmentation suggesting megaloblastic anaemia were seen in 26(26%) and 18(18%) patients respectively. On systemic examination haemic murmurs were detected among 22 patients (22%). Dashratham P et al, in their study found that 76% of patients had cardiac murmurs.¹² Hepatomegaly was the predominant finding on abdominal examination seen in 12 (12%) patients whereas palpable splenomegaly was seen in 7(7%) patients. Both liver and spleen were palpable in 7 patients. This study noted that 84% of cases presented as severe anaemia. This may be because of the reason that mild anaemia is neglected by people and they do not approach a doctor. Another reason may be illiteracy and lack of knowledge which makes them present to the hospital as severe anaemia cases. None of mild anaemia cases were noted in our study as they are often treated on outpatient basis and our study targeted inpatients. On peripheral smear examination microcytic hypochromic anaemia attributable to iron deficiency 50(50%) patients based on examination and observation of response to therapy was the most

common cause of anaemia . Similar findings were noted by Kouli R et al.¹⁶ This was followed by dimorphic anaemia was found in 27(27%) patients as the second most common cause of anaemia. Hence nutritional anaemia continues to predominate as the most common cause of anaemia in our part of world. Pure megaloblastic anaemia was seen in only 2 patient, 21 patients (21%) presented with normocytic normochromic anaemia .

CONCLUSION

Nutritional anaemia particularly iron deficiency anaemia is the most common cause of anaemia. It tends to affect the working age group and females predominantly. Patients continue to present with severe anaemia to the hospital.

REFERENCE

- Bernoist B, McLean E, Egli I, Cogswell M: Worldwide Prevalence of Anaemia 1993–2005: WHO Global Database on Anaemia. 2008, Geneva: World Health Organization Google Scholar
- Gardner B. Anemias of Inadequate Production. In: Kliegman RM, Behrman RE, Jenson H B, Stanton B F(eds). Nelson Textbook of Pediatrics. 18th edn. Philadelphia, W.B. Saunders. 2008:2011-4.
- Gupte S: Pediatric Hematology . In Gupte S. The Short Textbook of Pediatrics. 10th edn. New Delhi. Jaypee. 2004:454-62
- World Health Organization. The world health report 2002: reducing risks, promoting healthy life. World Health Organization; 2002.
- WHO, WHO global database on anaemia / Edited by Bruno de Benoist, Erin McLean, Ines Egli and Mary Cogswell, 2008
- Hussain R, Patwardhan VN. The influence of phytate on the absorption of iron. Indian J Med Res. 1959;47:676–682
- Hussain R. Dermal loss of iron in healthy Indian men. Ind Jour Med, 1960;48:235
- Azad KL, Dwivedi SK. A study of profile of patients with different degrees of anaemia. Int J Res Med Sci. 2017 Sep;5(9):4008-12.
- Alvarez-Uria G, Naik PK, Midde M, Yalla PS, Pakam R. Prevalence and severity of anaemia stratified by age and gender in rural India. Anaemia. 2014;2014.
- Talwelkar SR, Dhruva GA, Goswami SH. Survey of different types of anaemia. Schol J Appl Med Sci, 2019;7(12):3925-8.
- Anaemia ID, Assessment P. Control: A Guide for Programme Managers. World Health Organisation: Geneva: Switzerland; 2001.
- Dasharatham P, Reddy VS. A study of etiological and clinical profile of patients with severe anaemia in a tertiary care hospital. Int J Adv Med. 2018;5:1422-7.
- Gayathri BN, Rao KS. Pancytopenia: a clinico haematological study. J lab physic. 2011;3(1):15.
- Selvamuthukumaran S. A clinical study of severity of anaemia in adults attending outpatient department with regard to signs and symptoms in rural south indian medical college. Inter J Contempor Med Res. 2018;5(12):L21-L4.
- Unnikrishnan V, Dutta TK, Badhe BA, Bobby Z, Panigrahi AK. Clinico-aetiologic profile of macrocytic anaemias with special reference to megaloblastic anaemia. Ind J Hematol Blood Transfus. 2008 Dec 1;24(4):155-65.
- Kouli R, Pegu AK, Teronpi J, Shukla R, Bhattacharjee S, Sonowal S, et al. Study of anaemia in correlation with haematological parameters among adults in a tertiary care centre of North-East India. Int J Med Res Prof. 2016;2(4):84- 7.