

## Valuation of electrolyte status in tuberculosis and individuals

Sandeep Tripathi

Assistant Professor, Department of Biochemistry, National Institute of Medical Science & Research, Jaipur, Rajasthan, India

### ABSTRACT:

**Introduction:** This study is carried out with the objective to evaluate the serum electrolyte status among newly diagnosed cases of pulmonary TB which will be beneficial in preventing various complication and helpful for further appropriate treatment. **Materials and Methods:** Data such as name, age, gender etc. was recorded. All underwent clinical examination along with sputum smear examination by Z-N staining technique and chest X ray. Under aseptic conditions, 10 ml of venous blood samples of patients were collected and serum sodium, potassium and calcium levels were estimated using electrolyte analyzer which was based on principle of ion selective electrode. **Results:** Mean sodium level in group I was 133.2 mmol/L and in group II was 139.3 mmol/L, potassium level was 4.2 mEq/L in group I and 5.0 mEq/L in group II and calcium level was 5.3 mg/dl in group I and 8.9 mg/dl in group II. The difference was significant ( $P < 0.05$ ). **Conclusion:** Correction of these electrolyte imbalance could improve outcome of patient. Authors found that electrolytes were imbalanced in patients with tuberculosis as compared to healthy subjects. Thus, early diagnosis and prompt management of these abnormalities are critical.

**Keywords:** Electrolyte, tuberculosis, serum sodium, potassium, calcium

**Corresponding author:** Sandeep Tripathi, Assistant Professor, Department of Biochemistry, National Institute of Medical Science & Research, Jaipur, Rajasthan, India

**This article may be cited as:** Tripathi S. Valuation of electrolyte status in tuberculosis and individuals. J Adv Med Dent Scie Res 2015;3(2):267-269.

### INTRODUCTION

One third of the World population has been exposed to the TB bacterium, and new infections occur at a rate of one per second. In 2006, a total of 1.7 million people died of TB including 231,000 people with HIV.<sup>1</sup>

Tuberculosis typically affects all parts of the body especially lungs. The disease has become rare in developed countries, but is still a major public health problem in low- and middle-income countries.<sup>2</sup> For the past 40 years, treatment success in tuberculosis has been defined as the eradication of active infection whilst preventing resistance and recurrence, achieved through multidrug antimicrobial treatment.<sup>3</sup> In spite of newer modalities for diagnosis and treatment of TB, unfortunately, millions of people are still suffering and dying from this disease. TB remains one of the top 10 causes of death worldwide. Millions of people continue to fall sick from TB each year. India has the highest burden of both tuberculosis (TB) and multidrug-resistant (MDR) TB based on estimates reported in Global TB Report 2016. India accounts for about 24% of the global prevalence, 23% of the global incident cases, and 21% of the global.<sup>4</sup>

Electrolytes play a vital role in maintaining homeostasis within the body. Electrolyte imbalance can lead to impaired functions of heart, nervous system, muscular system, as well as lead to acid-base disorders. Decreased sodium electrolyte is the most common and frequent cause of electrolyte imbalance in all newly diagnosed tuberculosis patients.<sup>5</sup>

The lungs are the major site for Mycobacterium

tuberculosis primary infection and tuberculosis (TB) disease. Clinical manifestations of TB include primary TB, reactivation TB, laryngeal TB, endobronchial TB, lower lung field TB infection, and tuberculoma. Pulmonary complications of TB can include hemoptysis, pneumothorax, bronchiectasis, extensive pulmonary destruction, malignancy, and chronic pulmonary aspergillosis. Electrolytes play a vital role in maintaining homeostasis within the body. Electrolyte imbalance can lead to impaired functions of heart, nervous system, muscular system, as well as lead to acid-base disorders. Decreased sodium electrolyte is the most common and frequent cause of electrolyte imbalance in all newly diagnosed tuberculosis patients. TB can induce electrolyte imbalance mainly hyponatremia via several mechanisms containing local invasion to the adrenal glands (adrenal insufficiency), local invasion to hypothalamus or pituitary gland, tubercular meningitis and inappropriate ADH secretion via pulmonary infection, and excessive loss of these ions since diarrhea, vomiting, and sweating are frequently seen in PTB.<sup>6-8</sup> Hyponatremia is considered as one of the most common and important electrolyte abnormalities. The prevalence of hyponatremia was fifteen to thirty percent among tuberculosis patients. Similarly other electrolytes imbalance like hypochloremia and hypokalemia and reduced bicarbonate levels are also seen in pulmonary tuberculosis patients. Hypercalcemia have also been reported as one of the most common electrolyte imbalance in 25.7% of patients with TB. Hence this study is carried out with

the objective to evaluate the serum electrolyte status among newly diagnosed cases of pulmonary TB which will be beneficial in preventing various complication and helpful for further appropriate treatment.

## MATERIALS & METHODS

The present study was conducted among 90 newly diagnosed cases of pulmonary tuberculosis of both genders (Group I). Equal number of healthy subjects was included in group II. All were informed regarding the study and were involved after they agreed to participate.

Data such as name, age, gender etc. was recorded. All underwent clinical examination along with sputum

smear examination by Z-N staining technique and chest X ray. Under aseptic conditions, 10 ml of venous blood samples of patients were collected and serum sodium, potassium and calcium levels were estimated using electrolyte analyzer which was based on principle of ionselective electrode. Sodium concentration less than 136 mmol/L was termed as hyponatremia and less than 115 mmol/L as severe hyponatremia. Hypokalemia was defined as serum potassium levels <3.5 mEq/L. Hypocalcaemia was defined as serum calcium level < 4.5 mg/dl using flame emission spectrophotometry. Results thus obtained were subjected to statistical analysis. P value less than 0.05 was considered significant.

## RESULTS

Table 1 shows distribution of subjects in 2 groups.

**Table 1: Distribution of subjects**

Groups	Group I	Group II
Status	Tuberculosis	Healthy
M:F	50:40	50:40

Table 2, shows that mean sodium level in group I was 133.2 mmol/L and in group II was 139.3 mmol/L, potassium level was 4.2 mEq/L in group I and 5.0 mEq/L in group II and calcium level was 5.3 mg/dl in group I and 8.9 mg/dl in group II. The difference was significant ( $P < 0.05$ ).

**Table 2: Assessment of electrolytes**

Electrolytes	Group I	Group II	P value
Sodium	133.2	139.3	0.05
Potassium	4.2	5.0	0.05
Calcium	5.3	8.9	0.01

## DISCUSSION

In adults, tuberculosis is the second leading cause of death due to an infectious disease (after AIDS), with 95% of deaths occurring in low-income countries. The lungs are the major site for Mycobacterium tuberculosis primary infection and tuberculosis (TB) disease. Clinical manifestations of TB include primary TB, reactivation TB, laryngeal TB, endobronchial TB, lowerlung field TB infection, and tuberculoma.<sup>9</sup> Pulmonary complications of TB can include hemoptysis, pneumothorax, bronchiectasis, extensive pulmonary destruction, malignancy, and chronic pulmonary aspergillosis. Electrolytes play a vital role in maintaining homeostasis within the body. Electrolyte imbalance can lead to impaired functions of heart, nervous system, muscular system, as well as lead to acid-base disorders. Decreased sodium electrolyte is the most common and frequent cause of electrolyte imbalance in all newly diagnosed tuberculosis patients.<sup>10</sup> The present study was conducted to assess electrolyte status in tuberculosis and normal individuals.

In present study, mean sodium level in group I was 132.6 mmol/L and in group II was 138.2 mmol/L, potassium level was 3.5 mEq/L in group I and 4.0 mEq/L in group II and calcium level was 4.2 mg/dl in group I and 9.0 mg/dl in group II. A study by SS Warke et al, conducted on patients in India

evaluated the effect of treatment of antitubercular drugs on blood pH, electrolytes and osmolality, found mean value of serum Na<sup>+</sup> concentration found to be 134 mmol/L which was increased after treatment and reached to 143 mmol/L. Similarly Chloride level was increased 4 month posttreatment suggesting the decrease in reabsorptive capacity of uriniferous tubules towards chloride ions in tuberculosis.<sup>11</sup>

Olalekan et al<sup>12</sup> evaluated the effects of treatments on the imbalance of some electrolytes among patients infected with Mycobacterium tuberculosis. A total of one hundred and ten patients participated in this study. They were divided into four groups as follows: group 1 contains 50 normal patients without TB or HIV infection, group 2 20 new positive cases of TB patients without HIV infection (M = 13; F = 07), group 3 20 new positive cases of tuberculosis co-infected with HIV infection (M = 10; F = 10), and group 4 20 positive cases of TB patients on anti tuberculous drugs (M = 11; F = 09). Levels of sodium in TB patients on drugs (TBD) were significantly lowered compared to new case tuberculosis (NCT) patients ( $134.80 \pm 5.83$  mmol/L vs  $142.10 \pm 6.68$  mmol/L) while potassium levels were significantly elevated in TB patients on drugs compared with their new case counterparts ( $3.75 \pm 0.15$  mmol/L vs  $3.07 \pm 0.42$  mmol/L)  $P < 0.05$  respectively.

A study by SS Warke et al, conducted on patients in

India evaluated the effect of treatment of antitubercular drugs on blood pH, electrolytes and osmolality, found mean value of serum Na<sup>+</sup> concentration found to be 134 mmol/L which was increased after treatment and reached to 143 mmol/L. Similarly Chloride level was increased 4 month post treatment suggesting the decrease in reabsorptive capacity of uriniferous tubules towards chloride ions in tuberculosis.<sup>12</sup> Treatment of multidrug-resistant tuberculosis (MDR-TB) with second-line injectable drugs may result in an electrolyte imbalance. Electrolyte imbalances have been reported since the early use of capreomycin. In a study by Soeroto AY et al, they reported that After the first month of MDR-TB treatment, there was a significant decrease in mean serum potassium ( $4.0 \pm 0.4$  mEq/L to  $3.7 \pm 0.5$  mEq/L, ) in the kanamycin-based group and ( $4.1 \pm 0.5$  mEq/L to  $3.2 \pm 0.6$  mEq/L, ) in the capreomycin-based group. Serum potassium levels were significantly lower in the capreomycin-based group than in the kanamycin-based group ( $3.2 \pm 0.6$  mEq/L vs  $3.7 \pm 0.5$  mEq/L, ). Their findings emphasize the importance of routine monitoring of serum potassium, magnesium, and calcium during MDR-TB treatment, and that more attention should be paid when treatment is given using the capreomycin-based regimen. Moreover, their study supported the 2018 World Health Organization treatment guideline recommendations for removal of kanamycin and capreomycin from the MDR-TB regimens.<sup>13</sup> Similarly Amalia et al<sup>14</sup> and Rahmawati et al,<sup>15</sup> also demonstrated lower serum potassium levels in the capreomycin-based group than in kanamycin-based group.

## CONCLUSION

Screening for serum electrolytes in pulmonary tuberculosis is essential. Presence of coexisting metabolic disorder may cause severe health abnormalities and even death. Thus, early diagnosis and prompt management of these abnormalities are critical. Correction of these electrolyte imbalance could improve outcome of patient.

Authors found that electrolytes were imbalanced in patients with tuberculosis as compared to healthy subjects. Thus, early diagnosis and prompt management of these abnormalities are critical.

## REFERENCES

1. Lee P, Ho KK. Hyponatremia in pulmonary TB: evidence of ectopic antidiuretic hormone production. *Chest*. 2010;137 (1): 207–208.
2. Ganiger A, Patil L, Mrudula N. Evaluation of Serum Electrolyte Status among Normal Healthy Individuals and Newly Diagnosed Cases of Pulmonary TB in Tertiary Care Hospital in Bidar: An Observational Study. *Indian J Med Biochem* 2019;23(3):316–319.
3. Goli G, Mukka R, Sairi S. Study of serum electrolytes in acute exacerbation of chronic obstructive pulmonary disease patients. *Int J Res Med Sci* 2016; 4:3324–3327.
4. Ramos-Levi AM, Rodriguez-Hervada AD, Mendez-Bailon M, Macro-Marinez J. Drug induced hyponatremia: an updated review. *Minerva Endocrinol* 2014; 39:1–12.
5. Das P, Bandyopadhyay M, Baral K, Paul R, Banerjee AK. Dyselectrolytemia in chronic obstructive pulmonary diseases with acute exacerbation. *Nig J PhysiolSci* 2010; 25:25–2.
6. Jacobi J, Schnellhardt S, Kulschewski A, Amann KU, Kuefner MA, et al. An unusual case of hyponatraemia. *Nephrol Dial Transplant* 2010;25(3):998-1001.
7. Lee P, Ho KK. Hyponatremia in pulmonary TB: evidence of ectopic antidiuretic hormone production. *Chest*. 2010;137 (1): 207–208.
8. Ganiger A, Patil L, Mrudula N. Evaluation of Serum Electrolyte Status among Normal Healthy Individuals and Newly Diagnosed Cases of Pulmonary TB in Tertiary Care Hospital in Bidar: An Observational Study. *Indian J Med Biochem* 2019;23(3):316–319.
9. Terzano C, Di Stefano F, Conti V, Di Nicola M, Paone G, Peroianni A et al. Mixed acid-base disorders, hydro- electrolyte imbalance and lactate production in hypercapnic respiratory failure: the role of noninvasive ventilation. *PLoS One* 2012; 7:e35245.
10. Gopal Purohit, Sandeep Vernekar, C.R Choudhary, Sunil Vyas, Shameel Ahammed K K. Evaluation and comparison of serum electrolytes imbalance among patients with pulmonary tuberculosis, hiv, hiv coinfected with pulmonary tuberculosis & controls. *IJSR*. 2017; 6 (5):42-44.
11. K S. Warke, Z.H. Kban. Effect of anti-tuberculosis drug on serum electrolytes levels, osmolality, blood ph and pco<sub>2</sub> Levels in tuberculosis patients. *Asian J Microbial Biotech Env Sci* 2004; 6(1):89-91.
12. S. Warke, Z.H. Kban. Effect of anti-tuberculosis drug on serum electrolytes levels, osmolality, blood ph and pco<sub>2</sub> Levels in tuberculosis patients. *Asian J Microbial Biotech Env Sci* 2004; 6(1):89-91.
13. Soeroto AY, Darmawan G, Supriyadi R, Bhaskara PG, Santoso P, Alisjahbana B, Parwati I. Comparison of Serum Potassium, Magnesium, and Calcium Levels between Kanamycin and Capreomycin-BASEd Regimen-Treated MultiDrug-Resistant Tuberculosis Patients in Bandung (CEASE MDR-TB): A Retrospective Cohort Study. *International journal of microbiology*. 2019:1-6.
14. L. Amalia. Comparative study of kanamycin and capreomycin on serum potassium level of multidrug resistance tuberculosis patients at a hospital. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2016; 8(1):307–310.
15. R. N. Rahmawati, E. H. Pranggono, and R. Ruslami. Clinical characteristics and side effects of multidrug resistant tuberculosis therapy at top referral hospital West Java Indonesia. *Althea Medical Journal*. 2016; 3(4):526–532.