

ORIGINAL ARTICLE**Evaluate the relationship between pregnancy serum lipid concentrations and risk of preeclampsia**

Pankaj Bhushan

Assistant Professor, Department of Biochemistry, Hi-Tech Medical College and Hospital, Bhubaneswar, Odisha, India

ABSTRACT:

Aim: Evaluate the relationship between pregnancy serum lipid concentrations and risk of preeclampsia. **Material and Methods:** A cross sectional study was conducted in Department of Biochemistry, after taking the permission from the ethical committee. Total 300 patients include 100 non-pregnant, 100 pregnant and 100 PIH cases. Serum cholesterol, lipoproteins and triglyceride levels were evaluated for each of the three groups. **Results:** In the present study there was no significant difference in serum cholesterol levels between group I (normal, non-pregnant women) and group II (normal pregnant women). However, a significant difference was observed between group I and group III (pregnancy with PIH) the P values being <0.001. The serum cholesterol levels in group III were significantly higher than those in group I. However, there was a significant difference in serum cholesterol levels between group II and group III. The P value being <0.001. **Conclusion:** Pregnancy induced hypertension is a frequent complication during pregnancy which if advanced may be fatal for both mother and foetus. Endothelial damage might involve elevated serum triglycerides which may have a value that can be used as screening markers in early stages of pregnancy leading to preeclampsia in future.

Keywords: Lipid profile, PIH, pregnancy

Corresponding author: Pankaj Bhushan, Assistant Professor, Department of Biochemistry, Hi-Tech Medical College and Hospital, Bhubaneswar, Odisha, India

This article may be cited as: Bhushan P. Evaluate the relationship between pregnancy serum lipid concentrations and risk of preeclampsia. J Adv Med Dent Scie Res 2015;3(2):270-272.

INTRODUCTION

Hypertension is one of the most common medical problems encountered during pregnancy, complicating 2–3% of pregnancies. It has been reported that worldwide every minute a woman dies during labor or delivery. Maternal mortality rates are the highest in Africa, with a lifetime risk of 1 in 16; the rates are the lowest in western nations (1:2800), with a global ratio of 400 maternal deaths per 100,000 live births¹. Eclampsia accounts for 12% of such deaths². It is a common problem in developing countries, and its prevalence varies widely, from 1 in 100 to 1 in 1700³⁻⁶.

Gestational hypertension or pregnancy-induced hypertension (PIH) is the development of new hypertension in a pregnant woman after 20 weeks gestation without the presence of protein in the urine or other signs of preeclampsia.⁷ The alteration of serum lipid profile in essential hypertension is well documented. Serum lipids increase significantly during pregnancy and are further elevated twofold during pregnancy-induced hypertension (PIH)⁸. Disorders of lipoprotein metabolism are a major cause of endothelial dysfunction that may result in hypertension and proteinuria, clinical hallmarks of preeclampsia (PE)⁹. An abnormal lipid profile is known to be strongly associated with atherosclerotic cardiovascular diseases. The most important feature in toxemia of pregnancy is hypertension, which is supposed to be because of vasospastic phenomenon in kidney, uterus, placenta, and brain¹. The association

of serum lipid profile with gestational proteinuric hypertension is highly suggested to imply some new diagnostic tools. Moreover, the hormonal imbalance is a prime factor for the etiopathogenesis of PIH, and this endocrinal imbalance is well reflected in the alteration of serum lipid profile. Therefore, a simple measurement of serum lipid parameters may be of good predictive value in toxemia of pregnancy, avoiding the costly endocrinal investigations. In the literature, high serum lipoproteins have been reported in PIH.

Therefore, the present study was planned with the aim of assessment the Serum Lipoproteins in among non-pregnant women of reproductive age group 20-35 normal pregnant women without PIH and pregnant women with PIH of the same age group and to see whether there is any significant change in the above parameters in these groups.

MATERIAL AND METHODS

The present study was conducted in the Department of Biochemistry, After taking informed consent and the approval of the protocol review committee and institutional ethics committee.

INCLUSION CRITERIA

- Previously normotensive women with two repeat diastolic blood pressure measurements of ≥ 90 mmHg at third trimester of pregnancy
- Plus proteinuria of more than 300 mg/l in 24 hour or >2+ protein with dipstick.

EXCLUSION CRITERIA

- Women who were in labor,
- Women presenting with ruptured membranes,
- Women with multiple pregnancies,
- Women with any known concurrent medical complications
- Normal Pregnant women were those with diastolic blood pressure ≤ 90 mmHg at third trimester of pregnancy,
- Without any evidence of preeclampsia signs or proteinuria.

METHODOLOGY

Total 300 patients with a diagnosis of preeclampsia and eclampsia were selected for the study. Out of 300

RESULTS

Table 1: The estimation of serum cholesterol, triglycerides and lipoprotein fractions in nonpregnant women, normal pregnancy and pregnancy induced hypertension

Serum Lipids	Normal non pregnant women	Normal pregnancy	Pregnancy Induced Hypertension
Total serum cholesterol	210.57±13.65	220.06±17.98	235.87±34.74*
HDL Cholesterol	48.02±8.72	59.93±12.04*	46.24±7.21
LDL Cholesterol	139.20±15.72	116.70±29.07*)	136.20±30.78
VLDL Cholesterol	25.17±2.67	42.73±4.80*	54.13±6.37*
Triglycerides	121.04±10.87	224.89±22.27*	272.87±37.88*

Values are expressed as MEAN \pm SEM; n=5 P<0.001

DISCUSSION

Total Cholesterol In the present study, there was no significant alteration in total cholesterol levels could be observed in normal pregnancy as compared with normal non-pregnant women. These findings are similar to one of the findings of Sattar et al. However, the study of Sattar et al., also observed no significant alteration in pre-eclampsia¹⁰. In our study, the cholesterol level in group III (PIH) (235.87±34.74 mg/dl) are significantly higher than group I (210.57±13.65 mg/dl) and group II (220.06±17.98 mg/dl) . These findings are similar to that of Hubel CA, Who have found significant increase in serum total cholesterol in toxemia of pregnancy¹¹.

A significant increase in serum TG levels in third trimester of pregnancy was observed in the present study. Similar results were observed in previous studies Estrogen includes the hepatic biosynthesis of endogenous TG's which is carried by VLDL¹². In our study also serum TG levels were significantly higher in normal pregnant women (224.89±22.27 mg/dl) as compared with normal non-pregnant women 121.04±10.87mg/dl. Serum triglyceride concentrations rose significantly in PIH (272.87±37.88) as shown in our study which corroborated with the findings of many workers Jayantha De, Enquobahria, Cekmen^{13,14}. Increased Triglyceride, found in Pregnancy Induced Hypertension, is likely to be deposited in predisposed vessels.

A significant fall in LDL cholesterol levels in third trimester of normal pregnancy was also observed by other workers¹⁵. They have also reported a significant

increase in LDL in PIH. These findings are similar to our study. In our study LDL (136.20±30.78 mg/dl) fraction was significant in pregnancy induced hypertension subjects also when compared with non pregnant and normal pregnancy groups. In non-pregnancy LDL Cholesterol (139.20±15.72 mg/dl) was high than normal pregnancy (116.70±29.07 mg/dl) and PIH cases (136.20±30.78 mg/dl). Hypoestrogenaemia, predominance of smaller and denser serum LDL particles are supposed to be important contributors for endothelial dysfunction in PIH^{10,11,15}.

STATISTICAL ANALYSIS

The recorded data was compiled entered in a spreadsheet computer program and then exported to data editor page of SPSS version 25.0 (SPSS Inc., Chicago, Illinois, USA). Descriptive statistics included computation of percentages, means and standard deviation.

The increase in VLDL is due to hypertriglyceridemia leading to enhanced entry of VLDL that carries endogenous triglyceride into circulation. VLDL level further increased in PIH (54.13±6.37) as evidenced in the present study in corroboration with those of other worker^{16,17,10}.

In our study the HDL cholesterol were significantly increased in normal pregnancy 59.93±12.04 mg/dl over non-pregnant women (48.02±8.72 mg/dl) . We have also found a significant decrease in HDL-C in PIH (46.24±7.21 mg/dl) as compared with normal pregnant women. These findings correlate with the findings of Jayantha De¹³. The low level of HDL in preeclampsia is because of hypoestrogenemia and also due to insulin resistance¹⁸.

CONCLUSION

Pregnancy induced hypertension is a frequent complication during pregnancy which if advanced may be fatal for both mother and foetus. Endothelial

damage might involve elevated serum triglycerides which may have a value that can be used as screening markers in early stages of pregnancy leading to preeclampsia in future. Therefore, further studies are needed, to be done at multiple centers and higher numbers of cases to confirm our findings

REFERENCE

1. Nawal MN. An introduction to maternal mortality. *Rev Obstet Gynecol* 2008; 1:77–81.
2. World Health Organization (WHO). The World Health Report 2005: Make Every Mother and Child Count. Geneva, Switzerland: WHO, 2005, available online at: http://www.who.int/whr/2005/whr2005_en.pdf (accessed June 25, 2008).
3. Bagga R, Aggarwal N, Chopra V, Saha SC, Prasad GR, Dhaliwal LK. Pregnancy complicated by severe chronic hypertension: A 10-year analysis from a developing country. *Hypertens Pregnancy* 2007; 26:139–149.
4. Ayesha K, Nargi S. Eclampsia: An aggressive approach is needed. *Med Spectrum* 1998; 4:13–17.
5. Naseer D, Ataullah K, Nudrat E. Perinatal and maternal outcome of eclamptic patients admitted in Nishtar Hospital, Multan. *J Coll Physician Surg Pak* 2000; 10:261–264.
6. Bashir A, Aleem M, Shagufta J. Community education and downward trend in maternal deaths due to eclampsia. *Specialist Int* 1996; 12:147–154.
7. 40th. Williams obstetrics (24th ed.). McGraw-Hill Professional. 2014. ISBN 9780071798938
8. Uotila, JT, Tuimala RJ, Aarnio TM. Findings on lipid peroxidation and antioxidant function in hypertensive complication of pregnancy. *Br J Obstet Gynecol* 1993; 100:270–276.
9. Winkler K, Wetzka B, Hoffmann MM, et al. Triglyceride-rich lipoproteins are associated with hypertension in preeclampsia. *J Clin Endocrinol Metab* 2003; 88:1162–1166.
10. Sattar N, Bendoric A, Berry C, Shepherd J, Greer LA and Packard CJ. Lipoprotein subtraction concentration in preeclampsia: Pathogenic parallels to atherosclerosis. *Obst & Gynaecol*. 1997; 89 (3): 403-8.
11. Hubel CA, Lgcall F, Weissfeld L, Gandlay RE and Roberts JM. Small low-density lipoproteins and vascular cell adhesion molecule-1 are increased in association with hyperlipidemia in pre eclampsia. *metabolism*. 1998; 47 (10): 1281-8.
12. Glueck CJ, Fallet RU and Scheel D. Effects of oestrogenic compounds on triglyceride kinetics. *Metabolism*: 1975; 24: 537- 548.
13. Jayantha C, Ananda Kumar Mukhopadhyay and Pradip Kumar Saha *Indian Journal of Clinical Biochemistry*. Serum lipid profile in pregnancy induced hypertension. 2006; 21(2): 165-168
14. Eisenberg S and Levy RJ. Lipoprotein metabolism interrelationship between apoprotein and VLDL Enhanced LDL degradation and cholesterol synthesis. *Journal of Investigating Dermatology*. 1975; 101: 880-882.
15. Bary, Hunter and Herret, Wilde and Dakey. *Biochem Test for the Assessment of Feto Placental Function*. *Annals of Clin Bio*. 1979; 12: 83-107
16. . Kokia E, Barkai G, Reichman B, Segal P, Goldman B and Mashaoh S. Maternal serum lipid profile in pregnancies complicated by hypertensive disorders. *J. Perinat, Med (Germany)*. 1990; 18 (6): 473-8.
17. 23. Teichman AT, Wieland H, Cremer P, Kulow G and Mahle U. Serum Lipids in Pregnancies complicated by Hypertensive disorders. *Journal of Perinatal Medicine*. 1998; 22 (2): 107-111.
18. Mangal A, Shrivastava P, Gaur U, Jain A, Goyal U and Rath G. Department of Bio-chemistry, Anatomy & Obstet and Gynaecol, Analysis of placental alkaline phosphatase in hypertensive disorders complication pregnancy. *J Anat Soc India*. 2005; 54 (2) 1-9.