### Journal of Advanced Medical and Dental Sciences Research

@Society of Scientific Research and Studies

Journal home page: www.jamdsr.com

doi: 10.21276/jamdsr

(e) ISSN Online: 2321-9599;

(p) ISSN Print: 2348-6805

## Original Research

# Comparative study of heart rate variability in obese and normal young adults in Medical College

Vivek Saboo<sup>1</sup>, Sanjeev Kumar Sharma<sup>2</sup>

<sup>1</sup>Associate Professor, Department of Physical Medicine and Rehabilitation, National institute of <sup>2</sup>Medical Science and Research, Jaipur, India

Assistant Professor, Department of Physical Medicine and Rehabilitation, National institute of Medical Science and Research, Jaipur, India

#### **ABSTRACT:**

**Introduction:** Obesity is a condition from which the body has an excess accumulation of fat, which may cause from current daily routines and activities such as overeating and/or lacking of exercise. Nutritional problem in India is shifting from undernourishment to obesity. The autonomic nervous system plays an important role, not only in physiological situations, but also in various pathological settings such as diabetic neuropathy, myocardial infarction and congestive heart failure. **Materials and Method:** This study consists of 100 healthy young adults and 100 subjects in the age group of 20-26 years selected from the patient attendants and medical students of Santosh Medical College and Research Centre, Ghaziabad. Anthropometric parameters like height and weight were recorded. Body Mass Index (BMI) was calculated. They were categorized as Cases (n=100) and Controls (n=100) based on the BMI, Controls whose BMI <  $25 \text{kg/m}^2$  and Cases with BMI >  $30 \text{kg/m}^2$ . Subject was explained in detail about the ongoing procedure and ECG was digitally recorded after 10 minutes rest using lead II. **Results:** The mean age of the controls and cases were 22.16 and 21.82 respectively. Our study results showed significant inverse relationship between BMI and HRV parameters like SDNN, E/I ratio and HF nu, but at the same it showed significant positive relation of BMI and LF nu, LF/HF. **Conclusion:** The present findings demonstrate decreased HRV is associated with age, male gender and increased heart rate. **Keywords:** Obesity, Exercise, Nutritional, Heart Rate

Received: 27 February 2018

Revised: 19 March 2018

Accepted: 20 March 2018

Corresponding author: Dr. Sanjeev Kumar Sharma

**This article may be cited as:** Saboo V, Sharma SK. Comparative study of heart rate variability in obese and normal young adults in Medical College. J Adv Med Dent Scie Res 2018;6(5):90-94.

#### Introduction:

Obesity is a condition from which the body has an excess accumulation of fat, which may cause from current daily routines and activities such as overeating and/or lacking of exercise.<sup>1</sup> According to the International Association for the Study of Obesity (IASO), the International Obesity Task Force (IOTF) and the World Health Organization (WHO) proposed BMI cut-points 23.0 to 24.9 kg/m<sup>2</sup> for being overweight and above 25.0 kg/m<sup>2</sup> for obesity in adult Asians.<sup>2</sup>

Nutritional problem in India is shifting from undernourishment to obesity.<sup>3</sup> it is a condition, which has envolved with the advent of civilization, sedentary life style and high calorie diet.<sup>4</sup> Obesity is one of the causative factors for multiple co-morbid conditions leading to metabolic and cardiac disorders.<sup>5</sup> Growing number of evidences indicate association of obesity and sudden cardiac deaths.<sup>6,7</sup> Obesity is accompanied with varied combinations of abnormalities in the autonomic nervous system. One view is that obese people have a higher sympathetic tone that has been proved in some studies correlating with the

catecholamine levels. But there is also evidence of reduced cardiac sympathetic tone in some studies which was partly explained on the duration of obesity.<sup>8</sup>

The autonomic nervous system plays an important role, not only in physiological situations, but also in various pathological settings such as diabetic neuropathy, myocardial infarction and congestive heart failure. Autonomic imbalance with increased sympathetic activity and reduced vagal tone has been strongly implicated in the pathophysiology of development of arrhythmias and sudden cardiac deaths.<sup>9-11</sup> Among the different available non-invasive techniques for assessing the autonomic status, Heart Rate Variability (HRV) is a simple method to evaluate the sympathovagal balance at the sinoatriallevel.<sup>12</sup> Heart Rate Variability (HRV) is a specific and sensitive noninvasive tool to evaluate cardiac autonomic activity. HRV is the degree of variation of the heart rate under the balanced influence of sympathetic and parasympathetic components of the cardiac autonomic nervous system. HRV also indicates the extent of neuronal damage to autonomic nervous system.13

It is shown that in obese people, there is reduction in the HRV reflecting a predominant sympathetic influence on control of cardiac functions. HRV studies among adults and children with obesity have revealed inconsistent results including high and low sympathetic tones coupled with a reduction in vagal tone.<sup>14-16</sup> It has also been shown in western studies on obese adults that weight loss reverses back to parasympathetic control of cardiac functions.<sup>17</sup> While there are evidence of significant changes in autonomic control of cardiac functions in obese children and adults, there is lack of information concerning changes in obese young adults and also studies regarding HRV in obese Indian population are very few. Hence present study was under taken with the aim of evaluating the resting cardiac autonomic nerve activity given by the changes in HRV analysis in healthy obese young adults.

#### Materials and Method:

This study consists of 100 healthy young adults and 100 subjects in the age group of 20-26 years selected from the patient attendants and medical students of NIMS, Jaipur. Anthropometric parameters like height and weight were recorded. Body Mass Index (BMI) was calculated. They were categorized as Cases (n=100) and Controls (n=100) based on the BMI. Controls whose  $BMI < 25 \text{kg/m}^2$  and Cases with BMI $> 30 \text{kg/m}^2$ . Subject's clinical history and details were taken according to the standard proforma. Informed written consent was taken from all subjects in the study. Subjects with major illness like diabetes mellitus, hypertension, and endocrinal disorders and on any drugs affecting the Autonomic Nervous System like adrenergic blockers, calcium channel blockers, others were also excluded.

Subject was explained in detail about the ongoing procedure and ECG was digitally recorded after 10 minutes rest using lead II. First, a 5 minutes ECG in supine position was recorded, with subject breathing normally and used to determine the SDNN LF, HF and LF/HF. Then, the person was periodically instructed to take alternating 5 seconds of deep inspiration and 5 seconds of deep expiration for a period of 2 minutes and simultaneous ECG was recorded. This data was used to measure the E/I ratio. Heart Rate Variability software, was used in the computer, to detect the peak to peak intervals and further mathematical and analytical calculations in order to get the values of the parameters.

#### **Results:**

The mean age of the controls and cases were 22.16 and 21.82 respectively. Table-1 shows there was a statistically significant increase in BMI in cases when compared to controls. (p<0.001)

Variables	Control Mean	Cases Mean	'p' Value
Age (years)	22.16	22.94	0.158
BMI (Kg/m <sup>2</sup> )	24.34	31.36	< 0.001

Table-2 shows significantly reduced SDNN, E/I ratio, HF nu in cases when compared to controls suggesting the reduction in the parasympathetic activity. Significant increase in LF nu and LF/HF ratio shows elevated cardiac sympathetic activity in obese individuals. (p<0.01)

HRV	Control	Cases	'p' Value
	Mean	Mean	
SDNN(ms)	105.24	42.47	< 0.01
E/I ratio	1.76	1.40	< 0.01
LF <sub>nu</sub>	40.13	72.64	< 0.01
HF <sub>nu</sub>	61.38	29.13	< 0.01
LF/HF	0.68	3.26	< 0.01

Figure 1: comparison between control and cases



Our study results showed significant inverse relationship between BMI and HRV parameters like SDNN, E/I ratio and HF nu, but at the same it showed significant positive relation of BMI and LF nu, LF/HF.

#### **Discussion:**

This study is a novel part of the on-going long-term epidemiological Cardiovascular Risk in Young Finns study. Many studies into lifetime cardiovascular risk profiles in this study population have been reported previously. In 2003, Raitakari et al demonstrated that cardiometabolic risk profile assessed in 12-year to 18-year-old adolescents predicts adult carotid artery intima-media thickness (cIMT) independently of contemporaneous risk factors. These findings suggested that exposure to cardiometabolic risk factors early in life may induce changes in arteries that actually contribute to the development of atherosclerosis.<sup>18</sup>

Humans vary in their responsiveness to environmental stressors and HRV measurements may serve as physiological markers for such variation.<sup>19</sup> During normal sinus rhythm, the heart rate varies from beat to beat. Heart rate variability (HRV) results from the interplay between the various physiological mechanisms that regulate the heart rate. Since shortterm heart rate regulation is predominantly governed by sympathetic and parasympathetic neural activity, by examination of heart rate fluctuations normal functioning of the autonomic nervous system can be tested. HRV has been used as a measure of autonomic balance that emanates from endogenous sympathetic and parasympathetic rhythms which are partly modulated by respiratory sinus arrhythmia.<sup>20,21</sup> There are different measures of HRV- Time domain indices, Frequency domain indices and also non-linear techniques. Time domain analysis of heart rate variability uses statistical methods to quantify the variation of the standard deviation or the differences between successive RR intervals. Frequency domain analysis of heart rate variability enables us to calculate the respiratory dependent high frequency variations.<sup>12,22</sup>

This study was conducted to excess the effect on cardiac autonomic activity in healthy young audults 20-26 years. The major findings of this study indicate the presence of impaired par asympathetic activity and elevated level of sympathetic activity in obese group. When parasympathetic activity is considered there is not much difference among the studies, as almost all of the studies show a significant reduction in the parasympathetic activity with increasing body weight, which were similar to our studies.<sup>23-26</sup> In contrast to our study some studies showed a significant reduction in the sympathetic activity. This variation among the studies was partially explained on the basis of the duration of obesity.<sup>27</sup> It has been said that duration of the obesity has a major role to play in determining the level of cardiac sympathetic activity.<sup>28</sup> Correlation analysis between the obesity indices and HRV indices in this study showed that the time domain indices of HRV was significantly correlated negatively with overall obesity measured by BMI and WSR than the central obesity measured by WC, HC and WHR. In frequency domain indices LF component and LF/HF ratio were positively correlated whereas HF component was negatively correlated with both overall as well as central obesity indices. BFP was also negatively correlated with overall HRV and HF component which was not significant. There was also weak positive correlation between BFP and LF

component, LF/HF ratio. Age and sex matched regression analysis showed that BMI was the major determinant for the changes in both time domain and frequency domain indices. WSR was the next important predictor for the time domain indices whereas for the frequency domain indices it was WC and WHR.

Early detection and management by weight reduction and regular exercise can reduced the risk as these are shown to increase HRV.<sup>29,30</sup> HRV analysis can detect changes even before clinical signs appear.6 Thus regular assessment of HRV measures can be used as a biomarker for early detection and subsequent management of cardiovascular diseases in obese individuals.

This study is an effort to assess the effect of obesity on cardiac autonomic activity using Heart Rate Variability in young adults as sudden cardiac death in later ages can be prevented if life style modifications can be brought in earlier.

#### **Conclusion:**

The present findings demonstrate decreased HRV is associated with age, male gender and increased heart rate. Women have higher HF variability and lower LF variability than men., higher sympathetic and lower parasympathetic nerve activity in obese subjects. The results show that the HRV is decreased in overweight adults especially men indicating young sympathovagal imbalance. No changes were observed in HRV in underweight group. Changes in the autonomic nervous activity begin in the overweight and may become more prominent in the obese thus indicating increased cardiovascular risk. HRV indices were significantly associated with obesity indices. BMI was the major determinant for the changes in both time and frequency domain indices. Thus time and frequency domain analysis of the HRV in obese young adults shows imbalance in the autonomic neural activities to the heart

#### **References:**

- 1. H.A. Chethan, N. Murthy, and K. Basavaraju, "Comparative study of heart rate variability in normal and obese young adult males," Int J Biol Med Res, vol. 3, pp. 1621-1623, 2012.
- 2. The International Association for the Study of Obesity and the International Obesity Task Force. The Asia-Pacific perspective: redefining obesity and its treatment. Australia: IASO and IOTF, 2000
- Nageswari SK, Sharma R, Kohli DR. Assessment of respiratory and sympathetic cardiovascular parameters in obese school children. Indian J Physiol Pharmacol 2007; 51(3):235-43.
- Younghee K, Youn KS, Haymie C. MBI and metabolic disorders in South Korean adults: Korea National Health and Nutrition Survey. Obes Res 2004;12: 445-453.

- Yuko A, Hironobu Y, Mamoru K, Toshie S, Hiroshi E, Sukenobu I. VLCDInduced weight loss improves heart rate variability in moderately obese Japanese. Exp Biol Med 2001;226(5):440-5.
- 6. Kirsten LR, Harry H, Meena K, Eric B, Marek M, Michael M. Effects of moderate and vigorous physical activity on heart rate variability in british study of civil servants. Am J Epidemiol 2003;158:135-43.
- Frenco R, Bernard S, Andrea C, Tiziana G, Barbara DV, Ivana R et al. Assessment of cardiac autonomic modulation during adolescent obesity. Obes Rese 2003 April;11(4):541-548.
- 8. Jeong AK, Young GP, Kyung HC, Myung HH, Hee CH, Youn SC, Dokyung Y. Heart rate variability and obesity indices: Emphasis on the response to noise and standing. J Am Board Fam Pract 2005;18:97-103.
- 9. Vinik AI, Maser RE, Mitchell BD, Freeman R. Diabetic Autonomic Neuropathy. Diabetes Care 2003;26(5):1553–79.
- Kleiger RE, Miller JP, Bigger JT Jr, Moss AJ, The Multicenter Post-Infarction Research Group. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. Am J Cardiol 1987;59:256–62.
- 11. Bigger JT Jr, Fleiss JL, Steinman RC, Rolnitzky LM, Kleiger RE, Rottman JN. Frequency domain measures of heart period variability and mortality after myocardial infarction. Circulation 1992;85:164–71.
- Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Circulation 1996;93:1043–65.
- 13. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task force of the European society of cardiology and the North American society of pacing and electrophysiology. European Heart Journal 1996;17:354-81.
- Karason K, Molgaard H, Wikstrand J, Sjostrom L: Heart rate variability in obesity and the effect of weight loss. Am J Cardiol 1999;83:1242–7.
- 15. Gao YY, Lovejoy JC, Sparti A, Bray GA, Keys LK, Partington C: Autonomic activity assessed by heart rate spectral analysis varies with fat distribution in obese women. Obes Res 1996;4:55–63.
- Piccirillo G, Vetta F, Viola E, Santagada E, Ronzoni S, Cacciafesta M, et al: Heart rate and blood pressure variability in obese normotensive subjects. Int J Obes Relat Metab Disord 1998;22:741–50.
- 17. Akehi Y, Yoshimatsu H, Kurokawa M, Sakata T, Eto H, Ito S, et al. VLCD-Induced Weight

Loss Improves Heart Rate Variability in Moderately Obese Japanese. Exp Biol Med 2001;226:440–5.

- Raitakari OT, Juonala M, Kähönen M, Taittonen L, Laitinen T, Mäki-Torkko N, Järvisalo MJ, Uhari M, Jokinen E, Rönnemaa T, Åkerblom HK, Viikari JS. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. JAMA. 2003;290:2277-2283.
- 19. De Becker, P., Dendale, P., De Meirleir, K., et al (1998). Autonomic testing in patients with chronic fatigue syndrome. Am J Med. 105, :22S-26S.
- Hayano, J., Sakakibara, Y., Yamada, M., et al (1990). Decreased magnitude of heart rate spectral components in coronary artery disease. Its relation to angiographic severity. Circulation. 81, 1217-1224.
- Hrushesky, W.J (1991). Quantitative respiratory sinus arrhythmia analysis. A simple noninvasive, reimbursable measure of cardiac wellness and dysfunction. Ann NY Acad Sci. 618, 67-101.
- 22. Bilchick KC, Berger RD. Heart Rate Variability. J Cardiovasc Electrophysiol 2006;17(6):691–4.
- 23. Emdin M, Gastaldelli A, Muscelli E. Hyperinsulinemia and autonomic nervous system dysfunction in obesity: effects of weight loss. Circulation 2001; 103:513–9.
- 24. Martini G, Riva P, Rabbia F, Molini V, Ferrero GB, Cerutti F et al. Heart rate variability in childhood obesity. Clin Auton Res 2001 Apr; 11(2):87-91.
- 25. Facchini M, Malfatto G, Sala L, Silvestri G, Fontana P, Lafortuna C et al. changes of autonomic cardiac profile after a 3-week integrated body weight reduction program in severely obese patients. J Endocrinol Invest 2003 Feb;26(2):138-42.
- Poirier P, Hernandez TL, Weil KM, Shepard TJ, Eckel RH. Impact of dietinduced weight loss on the cardiac autonomic nervous system in severe obesity. Obes Res 2003 Sep; 11(9):1040-7.
- Masuo K, Mikami H, Ogihara T, Tuck ML. Weight gain-induced blood pressure elevation, Hypertension 2000;35:1135-40.
- Nagai N, Moritani T. Effect of physical activity on autonomic nervous system function in lean and obese children. Int J Obes Relat Metab Disord 2004 Jan;28(1):27
- 29. Akehi Y, Yoshimatsu H, Kurokawa M, Sakata T, Eto H, Ito S, et al. VLCD-Induced Weight Loss Improves Heart Rate Variability in Moderately Obese Japanese. Exp Biol Med 2001;226:440–5.
- Sandercock GR, Bromley PD, Brodie DA. Effects of Exercise on Heart Rate Variability: Inferences from MetaAnalysis. Med Sci Sports Exerc 2005;37(3):433–9.