

Original Research

Clinico-etiological profile of patients with acute pancreatitis

¹Nikita Sharma, ²Aguilera-Alvarez Victor H, ³Harsh Jitendra Kumar Patel, ⁴Alejandra Lopez, ⁵Saswat Kumar Jha, ⁶Sweta Sahu, ⁷Mandala Sri Shubha Reddy

¹MBBS, Government Medical College, Patiala, India;

²Imperial College London, Master of Public Health Country, Mexico;

³Intern, GMERS Medical College Gandhinagar, Gujarat, India;

⁴M.D, Mexico City, Universidad La Salle, Mexico;

⁵MBBS, Kathmandu Medical College, Teaching Hospital, Sinamangal, Kathmandu, Nepal;

⁶Second year MBBS Student, JJM Medical College, Davangere, Karnataka, India;

⁷MBBS, Dr. Pinnamaneni Siddhartha Institute of Medical Sciences and Research Foundation, Chinoutpalli, Vijayawada, Andhra Pradesh, India

ABSTRACT:

Aim: The present study was aimed to study etiology and complications of acute pancreatitis, to assess the clinical profile of acute pancreatitis. **Methods:** The study was a cross sectional study which was carried in the Department of General Surgery. Total 100 patients who were diagnosed for acute pancreatitis were include in this study and data collection on admission included age, sex, address and clinical presentation with respect to pain vomiting, gallstones trauma and drugs was noted. **Results:** Out of 100 patients included in study, 60 were males and 40 were females. In our study, majority of patients at the age group of 30-40 (41%) and followed by 40-50 years (32%). All the patients (100%) presented with pain abdomen, 83% of them presented with nausea/vomiting, 44% of them presented with fever and 27% of them with jaundice. In this study, 45% biliary pancreatitis was found to be the most common cause for acute pancreatitis. Alcoholism was the second most common cause (36%). Hyperlipidemia (3%) and traumatic (3%) pancreatitis was found in 3 patient each. Patients where no cause was found were labelled as idiopathic (13%). In males alcoholism induced pancreatitis was most common with a second commonest as biliary etiology. Diabetes mellitus was most prevalent in the study population 58%. Obesity as defined by the current definition was prevalent in 42%. **Conclusion:** Acute pancreatitis is one of the leading causes for increase morbidity and mortality to society. Clinical assessment along with lab markers correlated well with the mortality and morbidity.

Key Words: acute pancreatitis, clinical, morbidity, mortality.

Received: 25 September, 2021

Accepted: 28 October, 2021

Corresponding author: Nikita Sharma, MBBS, Government Medical College, Patiala, India

This article may be cited as: Sharma N, H Aguilera AV, Patel HJK, Lopez A, Jha SK, Sahu S, Reddy MSS. Clinico-etiological profile of patients with acute pancreatitis. J Adv Med Dent Scie Res 2021;9(11):102-105.

INTRODUCTION

The name "pancreas" is derived from Greek word "pan" (all) and "Kreas" (flesh). It was originally thought to act as cushion for stomach.¹ Acute Pancreatitis is an acute inflammatory process of the pancreas with variable involvement of regional tissues and remote organ system.² The average mortality rate in severe acute pancreatitis approaches 2-10%.³ Acute Pancreatitis is broadly classified (the Atlanta classification) as mild and severe: Mild acute pancreatitis is often referred to as interstitial pancreatitis, based on its radiographic appearance. Severe acute pancreatitis implies the presence of

organ failure, local complications, or pancreatic necrosis. Interstitial pancreatitis implies preservation of pancreatic blood supply; necrosis suggests the disruption of pancreatic blood supply with resulting ischemia. Most cases of acute pancreatitis fall into the mild category, with favorable recovery. However 15% to 20% cases of acute pancreatitis are severe and may result in a prolonged hospitalization, and local as well as systemic complications like systemic inflammatory response syndrome (SIRS), multi-organ system failure and death.⁴ With acute pancreatitis the inflammation comes on quickly over a few hours and usually goes away, leaving no permanently damage. However, it

can be fatal if complication occurs. There are many causes of acute pancreatitis, but the mechanisms by which these conditions trigger pancreatic inflammation have not been identified. Gallstone and alcohol abuses are the main causes of acute pancreatitis. The severity of Acute Pancreatitis can be predicted based upon clinically laboratory and radiological risk factors various severity grading system and serum markers. Some of this can be performed on admission to assist in triage of patient while others can be obtained during 1st 48 -72 hours or later. Severe acute pancreatitis is characterized by a short course, progressive MODS, early hypoxemia, increased incidence of necrosis, infection, and abdominal compartment syndrome (ACS).⁵ Multiorgan dysfunction syndrome, the extent of pancreatic necrosis, infection, and sepsis are the major determinants of mortality in Acute Pancreatitis.⁶ Pancreatic necrosis is considered as a potential risk for infection, which represents the primary cause of late mortality. Occurrence of acute respiratory (ARF), cardiovascular (CVF), and renal failures (RF) can predict the fatal outcome in SAP.⁷ Early accurate diagnosis is very important for its management. Symptoms of acute pancreatitis vary considerably. For this reason the clinician must carefully evaluate information derived from other sources that supplement the history and physical examination including laboratory tests, imaging studies before arriving at a correct diagnosis of acute pancreatitis. If the cause of the attack can be eliminated there will be no further attacks and the pancreas will return to normal in terms of its morphology and function.⁸ Management of acute pancreatitis has changed significantly over the past years. Early management is nonsurgical, solely supportive and patients with infected necrosis with worsening sepsis need intervention. Early intensive care has definitely improved the outcome of patients.⁹

MATERIAL AND METHODS

The study was a cross-sectional study which was carried in the Department of General Surgery, after

taking the approval of the protocol review committee and institutional ethics committee. All the 100 patients who were diagnosed for acute pancreatitis were included in this study. Patients with chronic pancreatitis, Renal failure, Cardiac failure and Generalized debility were excluded from this study.

The study population consisted of 100 cases of acute pancreatitis that fulfilled the diagnostic criteria. The diagnostic criteria included at least one of the three features. They are serum amylase more than 4 times the upper limit of normal, serum Lipase more than 2 times the upper limit of normal and ultrasound or CT scan suggestive of acute pancreatitis. This was based on the U. K. Guidelines for the management of acute pancreatitis. On admission history was collected and thorough physical examination was done. Data collection on admission included age, sex, address and clinical presentation with respect to pain vomiting, gallstones trauma and drugs was noted. History of previous episodes and co-morbidities was noted.

STATISTICAL ANALYSIS

The Descriptive procedure displays univariate summary statistics for several variables in a single table and calculates standardized values. Normally distributed continuous variables were expressed as mean (range) and non-normally distributed variables were expressed as median. Chi-square tests were employed to find out the difference between groups of frequencies obtained for the specific statements. A P value of less than 0.05 was considered statistically significant.

RESULTS

Out of 100 patients included in study, 60 were males and 40 were females. In our study, majority of patients at the age group of 30-40 (41%) and followed by 40-50 years (32%). The youngest patient was 16 year and the oldest Patient was 71 years (Table 1,2). All the patients (100%) presented with pain abdomen, 83% of them presented with nausea/vomiting, 44% of them presented with fever and 27% of them with jaundice (Table 3).

Table 1 Gender distribution of patients

Gender	Number	%
Male	60	60
Female	40	40

Table 2: Distribution of Age of acute pancreatitis patients

Age group in years	No of patients	%	P value
Below 20	3	3	0.81 NS
20-30	13	13	
30-40	41	41	
40-50	32	32	
50-60	8	8	
Above 60	2	2	

Table 3: Symptomatology of acute pancreatitis patients

Symptoms	No of patients	%
Pain abdomen	100	100
Fever	44	44
Vomiting	83	83
Jaundice	27	27

In this study, 45% biliary pancreatitis was found to be the most common cause for acute pancreatitis. Alcoholism was the second most common cause (36%). Hyperlipidemia (3%) and traumatic (3%) pancreatitis was found in 3 patient each. Patients where no cause was found were labelled as idiopathic (13%). In males alcoholism induced pancreatitis was most common with a second commonest as biliary etiology (Table 4).

Table 4: Etiology distribution of acute pancreatitis

Etiology	No of patients	%	P value
Biliary	45	45	0.07 NS
Alcoholism	36	36	
Hyperlipidaemia	3	3	
Traumatic	3	3	
Idiopathic	13	13	

Diabetes mellitus was most prevalent in the study population 58%. Obesity as defined by the current definition was prevalent in 42% (Table 5).

Table 5: Comorbidities in acute pancreatitis

Comorbidities	No of patients	%
Diabetes mellitus	58	58
Obesity	42	42

DISCUSSION

Acute pancreatitis is an inflammatory process of pancreas with varying involvement of regional tissues or remote organ system with potentially devastating consequences. Acute pancreatitis has a benign course, with gall stone and alcohol intake being cause of 70% of cases. Incidence of acute pancreatitis increases with age. Predicting the prognosis of a patient with acute pancreatitis at admission forms a very important strategy in management of Acute pancreatitis, considering this it enable us to practice guidelines for standardization of management of the patient which will in turn translate into improved outcomes.¹⁰ The present showed biliary pancreatitis (45%) as the most common cause for acute pancreatitis. The second most common cause was found to be alcoholism (36%). The other causes being hyperlipidaemia (3%), traumatic pancreatitis (3%) and 13% of patients did not show any symptoms and were labelled as idiopathic.

In biliary pancreatitis usually occurs in older adults, often have a history of cholelithiasis or intermittent, postprandial right upper-quadrant pain. Patients with acute pancreatitis present with mild to severe epigastric pain, with radiation to the back. Classically, the pain is characterized as constant, dull and boring, and is worse when the patient is supine.¹¹ The discomfort may lessen when the patient assumes a sitting or foetal position. A heavy meal or drinking binge often triggers the pain. In the present study all the patients (100%) presented with pain abdomen, 83% of them presented with nausea/vomiting, 44% of them presented with fever and 27 % of them with

jaundice. Vomiting may be severe and protracted. The abdominal distension was due to result of paralytic ileus arising from retroperitoneal irritation or ascites, or it may occur secondary to a retroperitoneal phlegmon. Jaundice may be occasionally seen in cases of gall stone pancreatitis, in which it represents distal CBD obstruction by gall stones.¹² Patients may also present with biliary colic.

On examination, severe pancreatitis was found to be associated with haemorrhage into the retro peritoneum may produce two distinctive sign's in about 3% of patients with pancreatitis namely Turner's sign (Bluish discoloration in the left flank) and Cullen's sign (Bluish discoloration of the periumbilical region).¹³ These are due to tracking of bloodstained retroperitoneal fluid through tissue planes of the abdominal wall to the flanks or along the falciform ligament. These signs suggest sever episode of acute haemorrhagic pancreatitis. A third rare finding called, fox sign (Bluish discoloration below the inguinal ligament or at the base of the penis) due to caudal tracking of fluid was also observed. Epigastric and right hypochondriac tenderness was present, sometimes present diffusely the abdomen. Bowel sounds were decreased or absent. Usually there were no masses palpable, if present it could be swollen pancreas or pseudocyst or abscess.

Temperature was mildly elevated (100-101 Degree F) even in uncomplicated cases. In severe cases, orthostatic hypotension and tachycardia may be present, along with tachypnea or even dyspnoea. There may be evidence of a pleural effusion, especially on the left side. In this study out of 100

patients, 60 were males and 40 were females. In our study, majority of patients at the age group of 30-40 (41%) and followed by 40-50 years (32%). The youngest patient was 16 year and the oldest Patient was 71 years. The age and sex-wise recruitment of the subjects in the present study was in accordance with the earlier studies.^{14,15} In other studies biliary pancreatitis was most prevalent. The combined etiology of alcohol and biliary pancreatitis is 81% which is fairly consistent with the other studies.¹⁶ In males alcoholism induced pancreatitis 50% was most common, second commonest is biliary etiology (40%). Kandasami P and colleagues reported that 78% of males the predominant etiology is alcoholism and 77% of females, the etiology for acute pancreatitis is biliary etiology.¹⁷ In the present study diabetes mellitus was most prevalent in the study population 58%. Obesity as defined by the current definition was prevalent in 42 %.

CONCLUSION

Acute pancreatitis is one of the leading causes for increase morbidity and mortality to society. Clinical assessment along with lab markers correlated well with the mortality and morbidity. Because pancreatitis mimics many other acute abdominal conditions, the diagnosis of acute pancreatitis must include a careful consideration of differential diagnosis, which should include perforated viscus, acute cholecystitis, appendicitis, and similar conditions.

REFERENCE

- Williams Norman, S. Bailey & Love's short practice surgery, 27th ed. 2018; 1212-39.
- Boon NA, College NR, Walker BR, Hunter AA. Davidson's principles and practice of medicine. London, Churchill Living stone, 20th ed. 2006; 904-9.
- Singh VK, Bollen TL and Wu Bu. An assessment of the severity of Interstitial pancreatitis. *Elm Gastroenterol Hepatol.* 2011; 9:1098.
- Stevens T, Parsi MA, Walsh RM. Acute pancreatitis: problems in adherence to guidelines. *Cleve Clin J Med.* 2009 Dec 1;76(12):697-704.
- Tao HQ, Zhang JX, Zou SC. Clinical characteristics and management of patients with early acute severe pancreatitis: experience from a medical center in China. *World journal of gastroenterology.* 2004 Mar 15;10(6):919.
- Buter A, Imrie CW, Carter CR, Evans S, McKay CJ. Dynamic nature of early organ dysfunction determines outcome in acute pancreatitis. *British Journal of Surgery.* 2002 Mar 1;89(3):298-302.
- Kong L, Santiago N, Han T Q, and Zhang S D, "Clinical characteristics and prognostic factors of severe acute pancreatitis," *World Journal of Gastroenterology.* 2004;10(22):3336-3338.
- Norton J, Berger G, Philip P (Editors) *Acute and Chronic pancreatitis; Harrison's Principles of Internal Medicine.* Newyork McGraw Hill. 20'h edition 2005; 1895-1906
- Chauhan S, Forsmark CE. The difficulty in predicting outcome in acute pancreatitis. *The American journal of gastroenterology.* 2010 Feb 1;105(2):443-5
- Elrnan R. Arneson N. Graham E: Value of blood amylase estimation in the diagnosis of pancreatic disease: A clinical study *Arch Surg.* 1929;19:943- 67.
- Glenn F, Frey C. Re-evaluation of the treatment of pancreatitis associated with biliary tract disease. *Ann Surg.* 1964;160:723-36.
- Ranson JH, Rifkmd KM, Roses DF. Prognostic signs and the role of operative management in acute pancreatitis. *Surg Gynecol Obstet.* 1974;139(1):69- 81.
- Acosta JM, Ledesma CL. Gallstone migration as a cause of acute pancreatitis. *N Engl J Med.* 1974;290(9):484-7.
- Werner J, Hartwig W, Uhl W, Müller C, Büchler MW. Useful markers for predicting severity and monitoring progression of acute pancreatitis. *Pancreatology.* 2003;3(2):115-27.
- Michael L. APACHE-II scores for assessment and monitoring of acute pancreatitis *The Lancet.* 1989;334(8656):201-5.
- Kandasami P, Harunarashid H, Kaur H. Acute Pancreatitis in a Multi-Ethnic Population Singapore *Med J.* 2002;43(6):284-88.
- De Waele JJ, Hoste EA, Malbrain ML. Decompressive laparotomy for abdominal compartment syndrome – a critical analysis. *Crit Care.* 2006;10:51-5