Journal of Advanced Medical and Dental Sciences Research

@Society of Scientific Research and Studies NLM ID: 101716117

Journal home page: www.jamdsr.com

doi: 10.21276/jamdsr

Index Copernicus value = 85.10

(e) ISSN Online: 2321-9599;

(p) ISSN Print: 2348-6805

Original Research

Knowledge of dental graduate students on management options for oral submucous fibrosis- An original research

Dr. Aishee Bhattacharya

BDS, Midnapore Medical College and Hospital, West Midnapore, West Bengal, India

ABSTRACT:

Aim: The purpose of the present study is to evaluate the knowledge of dental students regarding the aetiology as well as treatment options for Oral submucous fibrosis. **Methodology:** The questionnaire-based study was conducted among 100 participants. A self-administered questionnaire was used. The questionnaire-based study was conducted through an online forum, google form software. The questions were formed to observe the knowledge, attitude and practical approach of dental graduate students towards treating OSMF patients. The data collected were stored and results were analysed by SPSS software. **Result:** Out of 100 participants, 55% of participants reported that females have an increased prevalence of OSMF compared to males. The majority of participants were aware that betel quid was the most common habits and blanching mucosa were the common features associated with OSMF. 60% of participants were aware that stopping gutkha / pan chewing habits alone will not resolve the problem in patients having trismus. **Conclusion:** Students showed good knowledge on the various clinical and diagnostic aspects of Oral submucous fibrosis, however the knowledge on the management aspect of oral submucous fibrosis was moderate.

Keywords: Betel quid, blanching mucosa, malignancy, oral submucous fibrosis

Received: 25-04-2021 Revised: 05-05-2021 Accepted: 15-05-2021 Publ

Published: 30-05-2021

Corresponding author: Dr. Aishee Bhattacharya, BDS, Midnapore Medical College and Hospital, West Midnapore, West Bengal, India

This article may be cited as: Bhattacharya A. Knowledge of dental graduate students on management options for oral submucous fibrosis- An Original Research. J Adv Med Dent Scie Res 2021; 9(5):152-156.

INTRODUCTION

Oral submucous fibrosis (OSMF) is an oral precancerous condition characterized by inflammation and progressive fibrosis of the submucosal tissues resulting in marked rigidity and trismus. OSMF still remains a dilemma to the clinicians due to elusive pathogenesis and less well-defined classification systems. Pindborg has defined oral submucous fibrosis (OSMF) as an insidious chronic disease affecting any part of the oral cavity and occasionally extending to the pharynx, esophagus although occasionally preceded by and / or associated with Vesicle formation. OSMF is always associated with juxta-epithelial inflammatory reaction followed by fibroelastic changes in the lamina propria, with epithelial atrophy leading to stiffness of the oral mucosa causing trismus and difficulty in eating. The prevalence in India had increased in recent years to 6.42% with a higher predominance in the southern parts of the subcontinent. OSMF is seen commonly in males between 20 and 40 years of age. The common

sites involved are buccal mucosa, labial mucosa, retromolar pads, soft palate and floor of the mouth. Early features of OSMF include burning sensation, hypersalivation or xerostomia and blanching mucosa with marble-like appearance. Later mucosa becomes leathery and inelastic with palpable fibrous bands resulting in restricted mouth opening. Eventually, it leads to restriction of tongue movements, difficulty in swallowing, speech. Oral potentially malignant disorders (OPMDs) are considered as the early tissue changes that happen due to various habits such as smoking tobacco, chewing tobacco or stress.¹⁻⁵ The etiopathogenesis of OSMF is complex and it is a potentially malignant disorder attributed to areca nut (betel nut) chewing. The etiological factors include excessive chilli consumption, vitamins, iron deficiency, autoimmunity, genetic and environmental factors.⁶ Areca nut consists of alkaloids like arecoline, arecaidine, guvacine and guvacoline apart from flavonoids, tannins, catechin and copper. The alkaloids stimulate the fibroblasts to produce more

collagen, while its structure is stabilized by catechin and tannins.⁷ Apart from areca nuts, commercial products like gutkha, mawa and pan masala have been shown to cause Oral submucous fibrosis rapidly due to larger amounts of areca nut in these processed products and/or the synergistic action of nicotine over arecoline.⁸ gutkha products contain both SLT and areca nuts.⁹ The ideal goals of therapy of this potentially malignant disorder such as OSMF, include not only amelioration of the symptoms (burning sensation, restriction of mouth opening), but also stopping further disease progression and malignant transformation. Complete regression of this oral mucosal condition had not been achieved in any of the case studies reported till date hence an attempt at finding a permanent cure is still going on.¹⁰ Potentially malignant disorders (PMD) have a high risk of malignant transformation.¹¹ Malignancy is characterized by anaplasia, invasiveness, and metastasis.¹² Various treatment modalities including drug, surgical therapy and physiotherapy have been proposed till date for the management of OSMF. Surgical management of OSMF has involved excision of the fibrotic bands either by the scalpel or using a laser, with or without the use of interpositional grafts to maintain the oral opening. Various drugs with

antifibrotic, anti-inflammatory, and antioxidant activity¹³ have been used in the management of OSMF but with unpredictable results and incomplete remission. Not even a single drug has been reported to be effective in the treatment of OSMF. Hence, a combination of drugs has been used in the treatment of OSMF.¹⁴ The aim of the study was to assess the knowledge among dental students about the significance of Oral Submucous Fibrosis and its management.

AIM OF THE PRESENT STUDY

The purpose of the present study is to evaluate the knowledge of dental students regarding the aetiology as well as treatment options for Oral submucous fibrosis.

METHODOLOGY

The questionnaire-based study in Midnapore Medical College and Hospital, West Midnapore between 16.11.2020-16.4.2021, through online Google forms amongst 100 participants. Snowball sampling was used. A self-administered questionnaire was used which was in the English language and in an open-ended format.¹⁵ Questionnaire consisted of 13 questions. (Table1).

 Table 1- Questionnaire used in the present study

S.no.	Question				
1	Who is more affected by OSMF? (Male/ Female)				
2	Which is the most common habit associated with the incidence of OSMF?				
3	Most common clinical symptom noticed in patients?				
4	What are the various treatment modalities associated with all stages of OSMF?				
5	Do you think that oral cavity trismus will resolve if initiating factors (betel quid etc.) are stopped?				
6	Will oral physiotherapy alone will be sufficient to treat OSMF?				
7	Do you think OSMF has an aggressive malignant potential?				
8	Does hyaluronidase help with OSMF treatment?				
9	Which intralesional injection is used in the treatment of OSMF?				
10	Does treatment modalities vary with different stages of OSMF?				
11	Which one is the antioxidant amongst the drugs usually prescribed for OSMF?				
12	At what stage, intralesional injections are prescribed for treating OSMF?				
13	Which is the most common post-operative complication of OSMF?				

The questions were formed to assess the knowledge, attitude, practical approach of dentists towards managing and treating OSMF. The study protocol was approved by the Institutional Scientific Review board and ethical approval was also obtained. The results were analysed and the responses were tabulated in the form of a bar chart. Descriptive statistics for frequency and Chi square test were used to determine the correlation between the variables where P value < 0.05 is considered statistically significant.

RESULTS

From the results, we observed that 65% of participants were aware that females had an increased prevalence of OSMF compared to males. The majority of the participants responded that betel quid (53%) was the most common type of habit associated with OSMF. 66% of participants responded that blanching mucosa was the commonest feature associated with OSMF. 68% of participants were aware that OSMF had increased malignant potential and 32% of participants are unaware of it. 60% of participants were aware that

stopping of gutkha/ pan chewing habits alone will not resolve the problem in patients having trismus. 89% of participants were aware that treatment regimen / modalities vary with various stages of OSMF. 70% of participants were aware that they will treat OSMF by corticosteroids, lycopene, pentoxifylline. Majority of participants responded that antifibrinolytic activity will be the action of hyaluronidase in OSMF. 60% of participants were aware that drug lycopene was prescribed for OSMF as antioxidants. 53% of participants were aware that intralesional injections can be started from stage 3 of OSMF. 58% of participants were aware that corticosteroids + hyaluronidase / placental extracts will be the drugs/ drug given as intralesional injections in OSMF. 55% of participants were aware that oral physiotherapy alone will not help in treating OSMF. The majority of participants responded that oral mucositis will be the postoperative complication with surgical treatment of OSMF. The majority of respondents reported that OSMF has increased malignant potential (Pearson chi square test; p - value = 0.494 > 0.05 which was statistically not significant). Majority of respondents reported that OSMF treated corticosteroids, lycopene, pentoxifylline. (Pearson chi square test; p - value =

0.141 > 0.05, which was statistically not significant). The majority of respondents reported that treatment regimens / modalities vary with various stages of OSMF. (Pearson chi square test; p value = 0.413 > 0.05, which was statistically not significant). Majority of respondents reported that oral physiotherapy exercise alone cannot be helpful in treating OSMF (Pearson chi square test; p - value=0.988 > 0.05, which is statistically not significant). Majority of respondents reported drug lycopene prescribed for OSMF is antioxidants. (Pearson chi square; p -value = 0.042 > 0.05 which is statistically significant). (Table 2)

S. no.	Question	Answer received	P value
1	Who is more affected by OSMF? (Male/ Female)	35%- Male, 65%- Female	0.167
2	Which is the most common habit associated with the incidence of OSMF?	Betel quid (53%), Pan masala (33%), Tobacco (14%)	1.2
3	Most common clinical symptom noticed in patients?	Blanching (66%), xerostomia (22%), altered taste sensation (12%)	0.31
4	Do you know various treatment modalities associated with all stages of OSMF?	Yes (87%), No (13%)	0.029
5	Do you think that oral cavity trismus will resolve if initiating factor (betel quid etc.) are stopped?	Will not be resolved (60%), will be resolved (40%)	0.21
6	Will oral physiotherapy alone will be sufficient to treat OSMF?	Vill oral physiotherapy alone will be sufficient to treat OSMF?Will not help (55%), will help (45%)	
7	Do you think OSMF has an aggressive malignant potential?	More aggressive (68%), less aggressive (32%)	0.494
8	Does hyaluronidase help with OSMF treatment?	67% (helps), 33% (doesn't help)	0,199
9	Which intralesional injection is used in the treatment of OSMF?	Corticosteroid+hyaluronidase/ placenta extract (58%), others (42%)	0.141
10	Does treatment modalities vary with different stages of OSMF?	Yes (89%), No (11%)	0.413
11	Which one is the antioxidant amongst the drugs usually prescribed for OSMF?	Lycopene (74%), others (26%)	0.042
12	At what stage, intralesional injections are prescribed for treating OSMF?	Stage 3 (53%), Others (47%)	0.418
13	Which is the most common post- operative complication of OSMF?	78% (oral mucositis), scar (12%), necrosis (10%)	0,031

Table 2	2- Statistical	data	observed	in the	present	study
		-				

DISCUSSION

OSMF is an insidious, chronic disease with multifactorial etiology. Various treatment modalities had been proposed for OSMF but with unpredictable results. General dental practitioners have a basic knowledge about oral submucous fibrosis as they come across oral problems associated with OSMF in their practice. Generally, patients visit a dental clinic with a chief complaint associated with teeth pain or bleeding gums. Most of the time, practitioners give importance to the area of complaint and tend to miss out the asymptomatic mucosal lesions. If left unchecked, it can affect the quality of life and actually shorten life expectancy. This means that practitioners must play a more active role in educating their patients about the role of Oral submucous fibrosis and its malignant transformation and their overall health. An important component of health and overall quality of life is oral health. It can have an impact on the overall quality of life and dental diseases are increasing more and more in the world.¹⁶ From the results of this study, 68% of participants were aware

that OSMF had increased malignant potential. Murray C.J et al, reported similar evidence that OSMF had one of the higher rates of malignant potential among potentially malignant oral lesions. In patients with oral submucous fibrosis, the oral epithelium becomes atrophic and thereby becomes more vulnerable to carcinogens.¹⁷ 60 % of participants were aware that stopping gutkha / pan chewing habits alone will not resolve the problem in patients having trismus. Other papers also reported similar evidence that stopping the habits alone will not resolve the problem. OSMF does not regress spontaneously or on cessation of areca nut chewing. Once the disease is present, it either persists or becomes more severe with the involvement of additional areas of oral mucosa.¹⁸ In this study we observed that the majority of the participants responded that the action of hyaluronidase in OSMF will be antifibrinolytic activity. 86% of participants were aware that drug /drugs given for intralesional OSMF injection were corticosteroids in +hyaluronidase / placental extracts. Yadhav et al., reported the similar evidence that corticosteroids and hyaluronidase were the combinations of drugs used in intralesional injections.¹⁹ Leena et al reported that the medical treatments were not completely systematized, optimal doses of its treatment with intralesional injection of corticosteroids with hyaluronidase or placental extract are effective to some extent.²⁰ Borle and Borle postulated that treatment following intralesional injections of various drugs leads to aggravated fibrosis and pronounced trismus. Conservative lines of treatment like topical steroids, vitamins supplements, antioxidants, physiotherapy would give expected symptomatic relief of pain and burning sensation. The limitation of the study includes limited sample size, single centred study and does not represent ethnic groups or populations.

CONCLUSION

Students showed good knowledge on the various clinical and diagnosis aspects of Oral submucous fibrosis, however the knowledge on the management aspect of oral submucous fibrosis was moderate. A better knowledge of OSMF will endure safer health care services for the population.

REFERENCES

- Knowledge, Attitude, Practical Approach Of Dentist Towards Treating Oral Submucous Fibrosis Patients Attending Private Dental Hospital - A Survey Based Study. European Journal of Molecular & Clinical Medicine, 2020; 7(1):1221-1236.
- Chaitanya NC et al. 'Role of Vitamin E and Vitamin A in Oral Mucositis Induced by Cancer Chemo/Radiotherapy- A Meta-analysis', Journal of clinical and diagnostic research: JCDR. 2017; 11(5):ZE06–ZE09.
- Daga D et al. 'Efficacy of oral colchicine with intralesional hyaluronidase or triamcinolone acetonide in the Grade II oral submucous fibrosis', National journal of maxillofacial surgery. 2017; 8(1):50-54.

- Dharman S, Muthukrishnan A. 'Oral mucous membrane pemphigoid - Two case reports with varied clinical presentation', Journal of Indian Society of Periodontology. 2016; 20(6):630–634.
- 5. Hazarey VK et al. 'Oral submucous fibrosis: study of 1000 cases from central India', Journal of Oral Pathology & Medicine, 2006, 12-17p.
- Nigam, N. K. et al. (2014) 'Prevalence of oral submucous fibrosis among habitual gutkha and areca nut chewers in Moradabad district', Journal of oral biology and craniofacial research, 4(1), pp. 8–13.
- Trivedy, C. et al. (1997) 'Copper content in Areca catechu (betel nut) products and oral submucous fibrosis', The Lancet, p. 1447. doi: 10.1016/s0140-6736(97)24020-1.
- Karemore, T. V. and Karemore, V. A. (2011) 'Etiopathogenesis and Treatment Strategies of Oral Submucous Fibrosis', Journal of Indian Academy of Oral Medicine and Radiology, pp. 598–602. doi: 10.5005/jp-journals-10011-1230.
- Warnakulasuriya, S. and Muthukrishnan, A. (2018) 'Oral health consequences of smokeless tobacco use', Indian Journal of Medical Research, p. 35. doi: 10.4103/ijmr.ijmr_1793_17.
- Tilakaratne, W. M. et al. (2006) 'Oral submucous fibrosis: review on aetiology and pathogenesis', Oral oncology, 42(6), pp. 561–568.
- 11. Misra, S. R. et al. (2015) 'Metastatic hepatocellular carcinoma in the maxilla and mandible, an extremely rare presentation', Contemporary clinical dentistry, 6(Suppl 1), pp. S117–21.
- Subha, M. and Arvind, M. (2019) 'Role of Magnetic Resonance Imaging in Evaluation of Trigeminal Neuralgia with its Anatomical Correlation', Biomedical and Pharmacology Journal, pp. 289–296. doi: 10.13005/bpj/1640.
- Chaitanya, N. C. et al. (2018) 'An Insight and Update on the Analgesic Properties of Vitamin C', Journal of pharmacy & bioallied sciences, 10(3), pp. 119–125.
- 14. Daga, D. et al. (2017) 'Efficacy of oral colchicine with intralesional hyaluronidase or triamcinolone acetonide in the Grade II oral submucous fibrosis', National journal of maxillofacial surgery, 8(1), pp. 50–54.
- Knowledge, Attitude, Practical Approach Of Dentist Towards Treating Oral Submucous Fibrosis Patients Attending Private Dental Hospital - A Survey Based Study. European Journal of Molecular & Clinical Medicine, 2020; 7(1):1221-1236.
- Subashri, A. and Uma Maheshwari, T. N. (2016) 'Knowledge and attitude of oral hygiene practice among dental students', Research Journal of Pharmacy and Technology, p. 1840. doi: 10.5958/0974-360x.2016.00375.9.
- Pundir, S., Saxena, S. and Aggrawal, P. (2010) 'Oral submucous fibrosis a disease with malignant potential -Report of two Cases', Journal of Clinical and Experimental Dentistry, pp. e215–e218. doi: 10.4317/jced.2.e215.
- Nair, U. (2004) 'Alert for an epidemic of oral cancer due to use of the betel quid substitutes gutkha and pan masala: a review of agents and causative mechanisms', Mutagenesis, pp. 251–262. doi: 10.1093/mutage/geh036.
- Yadav, M. et al. (2014) 'Comparison of curcumin with intralesional steroid injections in Oral Submucous Fibrosis – A randomized, open-label interventional

Journal of Advanced Medical and Dental Sciences Research |Vol. 9|Issue 5| May 2021

study', Journal of Oral Biology and Craniofacial Research, pp. 169–173. doi: 10.1016/j.jobcr.2014.11.003.

20. Pindborg, J. J., Mehta, F. S. and Daftary, D. K. (1970) 'Occurrence of epithelial atypia in 51 Indian villagers with oral submucous fibrosis', British journal of cancer, 24(2), pp. 253–257.