

Original Research

Outcome of lateral splinterotomy in chronic anal fissure

Dr. Surendra Kumar

Assistant professor, Department of Surgery, Shri Vankateshwara Institute of Medical Sciences, Nh24, Venkateshwara Nagar, Near Rajabpur, Gajraula, Amroha, U.P.-244236, India

ABSTRACT:

Background: Lateral internal sphincterotomy (LIS) remains the gold-standard treatment of chronic anal fissure (CAF). The present study was conducted to assess outcome of lateral splinterotomy in chronic anal fissure. **Materials & Methods:** 65 patients underwent lateral splinterotomy in chronic anal fissure of both genders. Patients were recalled regularly and outcome was evaluated. **Results:** Out of 65 patients, males were 30 and females were 35. common complaints were pain during defecation in 92%, rectal bleeding in 84%, pruritis in 25%, constipation in 48% and perianal discharge in 14%. The common complications were rectal bleeding in 27%, perianal abscess in 4%, perianal hematoma in 3.7% and recurrence of lesion in 1.6%. Pain relief was seen at first week in 65%, at second week in 74%, at fourth week in 82% and at eighth week in 96% patients. The difference was significant ($P < 0.05$). **Conclusion:** LIS is the gold standard for the treatment of chronic anal fissure and patients found good pain relief. Common complication found was rectal bleeding.

Key words: Anal fissure, Lateral internal sphincterotomy, Pain

Received: 22 November 2020

Accepted: 24 December, 2020

Corresponding Author: Dr. Surendra Kumar, Assistant professor, Department of Surgery, Shri Vankateshwara Institute of Medical Sciences, Nh24, Venkateshwara Nagar, Near Rajabpur, Gajraula, Amroha, U.P.-244236, India

This article may be cited as: Kumar S. Outcome of lateral splinterotomy in chronic anal fissure. J Adv Med Dent Scie Res 2021;9(1):152-155.

INTRODUCTION

Anal fissure is a longitudinal tear in anoderm under the dentate line which mostly located posteriorly in the midline (90%). It is one of the most common benign diseases of anorectal area.¹ The main presenting symptom is pain during defecation, rectal bleeding, and emotional stress that it causes may reduce people's quality of life.² The healing of the chronic anal fissures takes longer than 8-12 weeks and in addition a hypertrophic papilla and a sentinel tubercle accompany the chronic anal fissure and the sphincter muscle fibers at the base of the tear are exposed.³

Lateral internal sphincterotomy (LIS) remains the gold-standard treatment of chronic anal fissure (CAF).⁴ LIS lowers the pressure exerted by the internal anal sphincter (IAS), restores normal perfusion of the anoderm, and leads to fast and durable pain relief and healing of the fissure, with a general complication rate of less than 7%.⁵ Recent innovations regarding nonsurgical alternatives have included the administration of topical nitrates (glyceryl trinitrate or isosorbide dinitrate) or botulinum toxin (Botox®, Allergan, Irvine, CA)

injections targeted to relax the IAS.⁶ Despite the initial excitement, randomized trials have revealed that topical glyceryl trinitrate is inferior to both Botox⁵ and LIS^{6,7} in providing symptomatic relief and fissure healing.⁷ The present study was conducted to assess outcome of lateral splinterotomy in chronic anal fissure.

MATERIALS & METHODS

The present study was conducted in general Surgery department comprised of 65 patients selected for lateral splinterotomy in chronic anal fissure of both genders. All were well informed and their informed written consent was obtained.

Demographic profile of all patients such as name, age, gender etc. was recorded. A thorough clinical examination of the site was carried out. Lateral splinterotomy (LIS) were carried out in the lithotomy position with open sphincterotomy under general or regional anesthesia. The anal canal was visualized with an anoscope, a longitudinal incision was made in the anoderm, and the distal half of the internal anal sphincter was divided under direct vision followed by closure of the mucosa.

All patients were asked to fill out a questionnaire that queried their symptoms. Anal pain was assessed before the treatment and at follow-up visits using a linear visual analog pain score. Anal incontinence was

assessed by means of a validated scoring and grading system, as reported by Pescatori et al. Patients were recalled regularly and outcome of treatment was recorded.

RESULTS

Table I Distribution of patients

Total- 65		
Gender	Males	Females
Number	30	35

Out of 65 patients, males were 30 and females were 35.

Table II Characteristics of patients

Complaint	Percentage	P value
Pain during defecation	92%	0.01
Rectal bleeding	84%	
Pruritis	25%	
Constipation	48%	
Perianal discharge	14%	

Table II, graph I shows that common complaints were pain during defecation in 92%, rectal bleeding in 84%, pruritis in 25%, constipation in 48% and perianal discharge in 14%. The difference was significant ($P < 0.05$).

Graph I Characteristics of patients





Table III Early and late complications outcomes

Complications	Percentage	P value
Rectal bleeding	27%	0.021
Perianal abscess	4%	
Perianal hematoma	3.7%	
Recurrence of lesion	1.6%	

Table III, graph II shows that common complications were rectal bleeding in 27%, perianal abscess in 4%, perianal hematoma in 3.7% and recurrence of lesion in 1.6%. The difference was significant ($P < 0.05$).

Graph II Early and late complications outcomes**Table IV Assessment of pain relief**

Duration	Percentage	P value
First week	65%	0.05
Second week	74%	
Fourth week	82%	
Eighth week	96%	

Table IV shows that pain relief was seen at first week in 65%, at second week in 74%, at fourth week in 82% and at eighth week in 96% patients. The difference was significant ($P < 0.05$).

DISCUSSION

The American Society of Colon and Rectal Surgeons (ASCRS) guidelines recommend initial nonsurgical management, which includes stool softeners, high fiber diet and warm sitz bath.⁸ However, a significant proportion of patients will fail conservative management, therefore further treatment options will be required.⁹ Lateral internal sphincterotomy (LIS) is attributed to be the gold standard for surgical management of chronic anal fissures when conservative and medical treatment fails. Besides its efficiency, LIS also have some risks of complications.¹⁰ Although incontinence, which is the most common and the feared one of those complications, was transient in most of the cases, 3% of the cases were considered to have it permanently at the end of the 72 months of follow up.¹¹ The present study was conducted to assess outcome of lateral splinterotomy in chronic anal fissure.

We found that out of 65 patients, males were 30 and females were 35. Common complaints were pain during defecation in 92%, rectal bleeding in 84%, pruritis in 25%, constipation in 48% and perianal discharge in 14%. Acar et al¹² evaluated the safe and adequate option of lateral internal sphincterotomy (LIS) in chronic anal fissure treatment. 417 patients who were treated for chronic anal fissure were

included. 228 (54.7%) were female and the mean age was 36.1 years (ranging from 17 to 73 years). Major complaints of patients; pain, bleeding, constipation, pruritus, perianal discharge. Recurrence occurred in 15 patients (3.6%) (12 males, three females) and eight patients (1.9%) developed incontinence (four with gas, four with soiling and seven females, one male). The complaints of all patients with gas incontinence and a patient with fluid incontinence regressed, whereas three patients had permanent fluid incontinency.

We found that common complications were rectal bleeding in 27%, perianal abscess in 4%, perianal hematoma in 3.7% and recurrence of lesion in 1.6%. Cadedu et al¹³ in their study 200 consecutive patients with chronic anal fissure, non responsive to previous treatment with nitroglycerin ointment or nifedipine, underwent LIS. Duration of operation, postoperative pain, duration of hospital stay, postoperative complications, time to resumption of work, recurrence and time to recurrence were assessed in all patients. The median operative time was 11 minutes (range 5–20); the median postoperative pain VAS score was 2 (range 0–4); every patient was mobilized on between 2 and 4 hours after surgery; the median hospital stay was 8 hours (range 7–10); the median time off work was 11 days (range 5–20 days). Neither impairment of

faecal incontinence nor relapse was detected in all patients at the end of observational period.

We found that pain relief was seen at first week in 65%, at second week in 74%, at fourth week in 82% and at eighth week in 96% patients. Montes et al¹⁴ in their study the fecal incontinence Quality of Life Scale was administered to any patient who had a Fecal Incontinence Severity Index score greater than 0 at 12 months postoperatively. The mean preoperative Gastrointestinal Quality of Life Index score was 118.34 T 6.33, which developed to 140.74 T 2.38 postoperatively ($P < 0.001$). At the two-month follow-up, 18 patients (7.38 percent) had a fecal Incontinence Severity Index score greater than 0. By 12 months, the number of patients with fecal Incontinence Severity Index score greater than 0 was reduced to seven (2.87 percent). These seven patients had a Gastrointestinal Quality of Life Index score similar to that of the group with postoperative fecal Incontinence Severity Index score of 0, and only three patients (1.22 percent) had evident deterioration in the Fecal Incontinence Quality of Life Scale. The 12-month total Gastrointestinal Quality of Life Index score of the three patients who developed anal abscess/fistula after sphincterotomy (139.33 T 3.21) was similar to the Gastrointestinal Quality.

CONCLUSION

Authors found that LIS is the gold standard for the treatment of chronic anal fissure and patients found good pain relief. Common complication found was rectal bleeding.

REFERENCES

1. Brady JT, Althans AR, Neupane R, et al. Treatment for anal fissure: is there a safe option? *Am J Surg.* 2017;214:623e628.
2. Collins EE, Lund JN. A review of chronic anal fissure management. *Tech Colorectal.* 2007;11:209e223.
3. Gil J, Luja'n J, Herna'ndez Q, Gil E, Salom MG, Parrilla P. Screening for the effectiveness of conservative treatment in chronic anal fissure patients using anorectal manometry. *Int J Colorectal Dis.* 2010;25:649e654.
4. Wald A, Bharucha AE, Cosman BC, Whitehead WE. ACG clinical guideline: management of benign anorectal disorders. *Am J Gastroenterol.* 2014;109:1141e1157.
5. Andjelkov K, Sforza M, Barisic G, Soldatovic I, Hiranyakas A, Krivokapic Z. Novel method for treatment of chronic anal fissure: adipose-derived regenerative cells e a pilot study. *Colorectal Dis.* 2017;19:570e575.
6. Acheson AG, Scholefield JH. Anal fissure: the changing management of a surgical condition. *Langenbeck's Arch Surg.* 2005; 390:1e7.
7. Yucel T, Gonullu" D, Oncu" M, Koksoy FN, Ozkan SG, Aycan O. Comparison of controlled-intermittent anal dilatation and lateral internal sphincterotomy in the treatment of chronic anal fissures: a prospective, randomized study. *Int J Surg.* 2009;7:228e231.
8. Perry WB, Dykes SL, Buie WD, Rafferty JF. Standards practice task force of the American Society of Colon and Rectal Surgeons Practice parameters for the management of anal fissures (3rd. revision). *Dis Colon Rectum.* 2010;53:1110-1115.
9. Nelson RL, Chattopadhyay A, Brooks W, Platt I, Paavana T, Earl S. Operative procedures for fissure in ano. *Cochrane Database Syst Rev.* 2011;9:CD002199.
10. Hsu TC, MacKeigan JM. Surgical treatment of chronic anal fissure. A retrospective study of 1753 cases. *Dis Colon Rectum.* 1984;27:475e478.
11. Montes BB, Tezcaner T, Yılmaz U, Leventoglu S, Oguz M. Results of lateral internal sphincterotomy for chronic anal fissure with particular reference to quality of life. *Dis Colon Rectum.* 2006;49:1045e1051.
12. Acar T, Acar N, Güngör F, Kamer E, Güngör H, Candan MS, Bağ H, Tarcan E, Dilek ON, Hacıyanlı M. Treatment of chronic anal fissure: Is open lateral internal sphincterotomy (LIS) a safe and adequate option?. *Asian journal of surgery.* 2019 May 1;42(5):628-33.
13. Cadeddu F, Selvaggio I, Muzi MG, Andreoli F, Amabile D, Milito G. Complications and results of lateral internal sphincterotomy for chronic anal fissure. *In BMC Geriatrics* 2009 Ap; 9 (1):1-1).
14. Menteş BB, Tezcaner T, Yılmaz U, Leventoglu S, Oguz M. Results of lateral internal sphincterotomy for chronic anal fissure with particular reference to quality of life. *Diseases of the colon & rectum.* 2006 Jul;49(7):1045-51.