Journal of Advanced Medical and Dental Sciences Research

@Society of Scientific Research and Studies NLM ID: 101716117

Journal home page: www.jamdsr.com

doi: 10.21276/jamdsr

Index Copernicus value = 85.10

(e) ISSN Online: 2321-9599;

(p) ISSN Print: 2348-6805

Original Research

Incidence of Transalveolar Extraction in Upper First Molar in patients Visiting Department of Oral surgery Dental college

Nahida Dar

Oral and Maxillofacial Surgeon, GDC, Srinagar, Jammu and Kashmir, India

ABSTRACT:

Transalveolar extraction is a method employed when normal forceps extraction is notpossible due to various difficulties which included tooth fracture during normal extractions. Though this technique is employed in clinical practice, it can lead to several complications aftersurgery. Hence, the objective of this study was to evaluate the incidence of transalveolarextractions of upper first molar. A retrospective study was conducted. Data collection was donethrough reviewing records of 500 patients visiting a exodontia unit inoral surgery gdc Srinagar from June(2019) toMarch(2020). A total of 500 patients who had undergone extraction of upper first molar were evaluated. Data such as age, gender, type of extraction and tooth number were evaluated. Bothfrequency and Chi-square tests were done through IBM SPSS statistical analysis. Prevalence oftransalveolar extractions in the upper first molar was 1.5% in the study population. Among theindividuals who had undergone transalveolar extraction, 59.09% were males and 40.91% werefemales. About 59.09% of the transalveolar extractions was 1.5% of the upper first molar and more prevalent among the 20-40 age group.

Keywords: defense Incidence, Surgical approach, Tooth fracture, Transalveolar extraction, Upper first molar, Innovative technique

Received: 18 November, 2020

Accepted: 20 December, 2020

Corresponding author: Nahida Dar, Oral and Maxillofacial Surgeon, GDC, Srinagar, Jammu and Kashmir, India

This article may be cited as: Dar N. Incidence of Transalveolar Extraction in Upper First Molar in patients Visiting Department of Oral surgery Dental college. J Adv Med Dent Scie Res 2021;9(1):171-175.

INTRODUCTION

Tooth extraction is a most commonly performed procedure in developing countries(Saheeb and Sede, 2013). An ideal extraction of tooth is considered as painless removal of the whole tooth, or a tooth root with minimal trauma to the investing tissues, so that the uneventful healing of the wound occurs(Narasimman, 2018). Extraction of a tooth is performed for several reasons including dental caries. periodontal diseases, orthodontic reasons, impacted teeth, failed dental treatment, prosthetic indications and other reasons. From the cross- sectional studies conducted across different countries, dental caries and periodontal diseases are the major cause of the tooth loss(Agerholm and Sidi, 1988; Corbet and Davies, 1991; Haddad et al., 1999). The maxillary first molar can be a very difficult tooth to extract atraumatically. The root morphology contains a very divergent root pattern with a thick palatal root. It is very adjacent to the maxillary sinus which is the most common cause of oro-antralcommunication(Polat, Ay and Kara, 2007). A study by Shah et al(Shah, Faldu

and Chowdhury, 2019) reported the first molar to be the most frequently extracted teeth. This is due to several reasons including as the first permanent tooth to be erupted in the oral cavity and broader surface area with pit and fissures making them more susceptible to plaque accumulation and caries formation Extraction of teeth requires the combination of principles of surgery and elementary physical mechanisms. When these principles are applied correctly the tooth can be removed from the socket without any use of a large amount of strength and force. Excessive force without a proper surgical can damage the surrounding bone and teeth. Such excessive force occasionally damages or fractures the leading the need of surgical crown to extraction/transalveolar extraction. Transalveolar extractions are the surgical removal of teeth by drilling the bone after the elevation of the flap. It is one of the most commonly performed minor oral surgical procedures in maxillofacial surgery for theimpacted third molar removal(Bui, Seldin and Dodson, 2003). However, it can cause various post

operative complications including dry socket, pain, trismus and swelling(Grossi et al., 2007).

Our department is passionate about research we have published numerous high quality articles in this domain over the past years ((Kavitha et al., 2014), (Praveen et al., 2001),(Devi and Gnanavel, 2014), (Putchala et al., 2013), (Vijayakumar et al., 2010), (Lekha et al., 2014a, 2014b) (Danda, 2010) (Danda, 2010) (Parthasarathy et al., 2016) (Gopalakannan, Senthilvelan and Ranganathan, 2012), (Rajendran et al., 2019), (Govindaraju, Neelakantan and Gutmann, 2017), (P. Neelakantan et al., 2015), (PradeepKumar et al., 2016), (Sajan et al., 2011), (Lekha et al., 2014a), (Neelakantan, Grotra and Sharma, 2013), (Patil et al., 2017), (Jeevanandan and Govindaraiu, 2018), (Abdul Wahab et al., 2017), (Eapen, Baig and Avinash, 2017), (Menon et al., 2018), (Wahab et al., 2018), (Vishnu Prasad et al., 2018), (Uthrakumar et al., 2010), (Ashok, Ajith and Sivanesan, 2017), (PrasannaNeelakantan et al., 2015). Now we are focussing on epidemiological studies. There are no studies done regarding the incidences and prevalence of transalveolar extractions. Hence in the present study, we evaluate the incidence of transalveolar extractions of upper first molars among the patients visiting a dental hospital.

MATERIALS AND METHODS

This study was conducted in a Dental College Oral Surgery Department. The study samples were chosen from the patients visiting a hospital in SRINAGAR from June(2018) to March(2020).

DATA COLLECTION

A retrospective study was carried out on patients of all ages who had undergone extraction of 16 and 26. Data collection was done through reviewing the records of 1000 between June(2019) - March(2020). Data such as age, gender, type of extraction, tooth number were noted. The data collected was verified with intraoral photographs.

INCLUSION CRITERIA

Patients of all groups who had undergone extraction of 16 and 26 were included.

EXCLUSION CRITERIA

Patient who are under special care, dentofacial trauma were excluded from this study Incomplete/censored data were excluded too.

DATA ANALYSIS

The data collected was entered in an Excel sheet. Data was analysed through frequency tests and cross tabulations using SPSS software.

RESULTS AND DISCUSSION

A total of 500 patients were reported to have undergone extraction of 16 and 26 in the span of 9 months(June(2019) - March(2020)). Among them, 1.51%(22) of the patients had undergone transalveolar extractions of maxillary first molar. In the study population 54.8% were males and 45.1% were females. The incidence of transalveolar extraction was higher in males(59.9%) than females(40.9%), however which is statistically insignificant(Chi square; p>0.05)

Majority of the patients who had undergone transalveolar extractions belong to the age group of 20-40 years(59.9%) and about 31.82% of the patients belong to the age group of 40-60 years. There was no difference in the tooth number in which had undergone transalveolar extractions(Maxillary right first molar(16) - 50%, Maxillary left first molar(26) - 50%).

Though exodontia is the most common surgical procedure performed in the maxillofacial surgery, its complications are unexpected events that can lead to increased morbidity which occasionally can progress to prolonged phase of treatment(Venkateshwaret al., 2011) SEP Dental caries are the most common infectious diseases in the world and about 60%-90% of the students are affected by it(Saber et al., 2018). Cahen et al(Cahen, Frank and Turlot, 1985) reported various reasons for tooth extraction such as dental caries. endodontic problems such as pulp inflammation, necrosis or tooth fracture, periodontal diseases, eruption problems, extractions for prosthetic orthodontic treatment, and trauma, occlusal dysfunctions such as extrusion, and iatrogenic factors. The maxillary first molar is one of the permanent teeth which erupts early and is more prone to carious exposure and it is important in maintaining normal masticatory function(ADA Division of Communications. Journal of the American Dental Association and ADA Council on Scientific Affairs. 2006).

In the present study, there were about 500 patients who had undergone extraction of maxillary first molar. Previous studies by Ozmen et al(Ozmen, 2019) and George et al(George et al., 2011) reported that the maxillary first molar was the most frequently extracted tooth. In the present study, there was no significant difference between genders associated with tooth loss(males - 54.8%, females - 45.4%). However few studies reported higher prevalence of maxillary first molar tooth loss among males(Barbato and Peres, 2009; Jafarian and Etebarian, 2013). This may be due to the higher aesthetic concerns and awareness among females.

Extractions could lead to various complications including haemorrhage, persistent pain, swelling, infections, dry socket, dentoalveolar fractures, oral sinus communications etc,. These are influenced by various factors including age, health condition, tooth level, surgeon's experience, smoking, contraceptive medications, surgical technique(Aravinth and Ganapathy, 2019). Fracture of maxillary tuberosity is one of the most common complications in the extraction of the upper first molar(Altuğ et al., 2009).

Baniwal et al(Baniwalet al., 2007) studied the complications of exodontia in tertiary centres and reported that the most frequent complication was the fracture of tuberosity. Controlled force and proper surgical planning is required to avoid the fracture of bone or tooth crown during extractions.

In the present study, the incidence of transalveolar extraction of the upper first molar was 1.51% and occurred more commonly in the age group of 20-40 years. Transalveolar extraction can lead to various complications than normal extractions. A study by Adwani et al(Adwaniet al., 2012) reported a higher incidence of dry socket in transalveolar extraction than intra alveolar extraction. Postoperative complications from transalveolar extractions occur at the rate ranging from 10% to 12% (Burke, 1961; Natarajan, Balakrishnan and Thangavelu, 2017).

Incidence of the transalveolar extraction and other complications can be limited by giving meticulous importance to surgical details including patient's preparation, asepsis, proper management of soft and hard tissues, controlled force of surgical instruments etc,.

CONCLUSION

Within the limitations of this study, the incidence of transalveolar extractions of the upper first molar was found to be 1.51%. It was more common in the age group of 20-40 years. Since there are no studies conducted on the incidence of transalveolar extractions, more epidemiological studies with large samples should be conducted to rule out the definite cause and outcome of those surgical approaches.

REFERENCES

- 1. Abdul Wahab, P. U. et al. (2017) 'Risk Factors for Post-operative Infection Following Single Piece Osteotomy', Journal of maxillofacial and oral surgery, 16(3), pp. 328–332.
- 2. Abhinav, R. P. et al. (2019) 'The Patterns and Etiology of Maxillofacial Trauma in South India', Annals of maxillofacial surgery, 9(1), pp. 114–117.
- 3. ADA Division of Communications, Journal of the American Dental Association and ADA Council on Scientific Affairs (2006) 'For the dental patient. Tooth eruption: the permanent teeth', Journal of the American Dental Association, 137(1), p. 127.
- Adwani, N. et al. (2012) 'Incidence of dry socket following intraalveolar and transalveolar extraction: A comparative study', Journal of DattaMeghe Institute of Medical Sciences University, 7(4), pp. 228–233.
- Agerholm, D. M. and Sidi, A. D. (1988) 'Reasons given for extraction of permanent teeth by general dental practitioners in England and Wales', British dental journal, 164(11), pp. 345–348.
- 6. Aravinth, H. and Ganapathy, D. (2019) 'Knowledge about post-extraction complications among dental students', Drug Invention Today.
- Ashok, B. S., Ajith, T. A. and Sivanesan, S. (2017) 'Hypoxia-inducible factors as neuroprotective agent in Alzheimer's disease', Clinical and experimental pharmacology & physiology, 44(3), pp. 327–334.

- Awasthi, M. (2018) 'Basic Principles of Oral and Maxillofacial Surgery', Manual for Dental Hygienist, pp. 284–284. doi: 10.5005/jp/books/14199_54.
- 9. Baniwal, S. et al. (2007) 'Prevalence of complications of simple tooth extractions and its comparison between a tertiary center and peripheral centers: a study conducted over 8,455 tooth extractions', JNMA; journal of the Nepal Medical Association, 46(165), pp. 20–24.
- 10. Barbato, P. R. and Peres, M. A. (2009) 'Tooth loss and associated factors in adolescents: a Brazilian population-based oral health survey', Revista de saudepublica, 43(1), pp. 13–25.
- Bui, C. H., Seldin, E. B. and Dodson, T. B. (2003) 'Types, frequencies, and risk factors for complications after third molar extraction', Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons, 61(12), pp. 1379–1389.
- 12. Burke, J. F. (1961) 'The effective period of preventive antibiotic action in experimental incisions and dermal lesions', Surgery, 50, pp. 161–168.
- Cahen, P. M., Frank, R. M. and Turlot, J. C. (1985) 'A survey of the reasons for dental extractions in France', Journal of dental research, 64(8), pp. 1087–1093.
- Christabel, A. et al. (2016) 'Comparison of pterygomaxillarydysjunction with tuberosity separation in isolated Le Fort I osteotomies: a prospective, multi-centre, triple-blind, randomized controlled trial', International Journal of Oral and Maxillofacial Surgery, pp. 180–185. doi: 10.1016/j.ijom.2015.07.021.
- Corbet, E. F. and Davies, W. I. (1991) 'Reasons given for tooth extraction in Hong Kong', Community dental health, 8(2), pp. 121–130.
- 16. Danda, A. K. (2010) 'Comparison of a single noncompressionminiplate versus 2 noncompressionminiplates in the treatment of mandibular angle fractures: a prospective, randomized clinical trial', Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons, 68(7), pp. 1565– 1567.
- Devi, V. S. and Gnanavel, B. K. (2014) 'Properties of Concrete Manufactured Using Steel Slag', Procedia Engineering, 97, pp. 95–104.
- Eapen, B. V., Baig, M. F. and Avinash, S. (2017) 'An Assessment of the Incidence of Prolonged Postoperative Bleeding After Dental Extraction Among Patients on Uninterrupted Low Dose Aspirin Therapy and to Evaluate the Need to Stop Such Medication Prior to Dental Extractions', Journal of maxillofacial and oral surgery, 16(1), pp. 48–52.
- George, B. et al. (2011) 'Prevalence of permanent tooth loss among children and adults in a suburban area of Chennai', Indian journal of dental research: official publication of Indian Society for Dental Research, 22(2), p. 364.
- Gopalakannan, S., Senthilvelan, T. and Ranganathan, S. (2012) 'Modeling and Optimization of EDM Process Parameters on Machining of Al 7075-B4C MMC Using RSM', Procedia Engineering, 38, pp. 685–690.
- Govindaraju, L., Neelakantan, P. and Gutmann, J. L. (2017) 'Effect of root canal irrigating solutions on the

compressive strength of tricalcium silicate cements', Clinical oral investigations, 21(2), pp. 567–571.

- 22. Grossi, G. B. et al. (2007) 'Assessing postoperative discomfort after third molar surgery: a prospective study', Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons, 65(5), pp. 901–917.
- Haddad, I. et al. (1999) 'Reasons for extraction of permanent teeth in Jordan', International dental journal, 49(6), pp. 343–346.
- 24. Jafarian, M. and Etebarian, A. (2013) 'Reasons for extraction of permanent teeth in general dental practices in Tehran, Iran', Medical principles and practice: international journal of the Kuwait University, Health Science Centre, 22(3), pp. 239– 244.
- Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', European Archives of Paediatric Dentistry, pp. 273–278. doi: 10.1007/s40368-018-0356-6.
- 26. Jesudasan, J. S., Wahab, P. U. A. and Sekhar, M. R. M. (2015) 'Effectiveness of 0.2% chlorhexidine gel and a eugenol-based paste on postoperative alveolar osteitis in patients having third molars extracted: a randomised controlled clinical trial', The British journal of oral & maxillofacial surgery, 53(9), pp. 826– Journal of Contemporary Issues in Business and Government | Vol 27, Issue 2, 2021 347
- 27. OVIYA. V. J et al / Incidence of Transalveolar Extraction in Upper First Molar Among Patients Visiting A Dental Hospital 830.
- Katoma, M., Siziya, S. and Sichilima, A. (2017) 'The most frequently extracted teeth type: a retrospective cross-section study at Ndola teaching hospital, low cost dental clinic, Zambia', Asian Pacific Journal of Health Sciences.
- Kavitha, M. et al. (2014) 'Solution combustion synthesis and characterization of strontium substituted hydroxyapatite nanocrystals', Powder Technology, 253, pp. 129–137.
- Kumar, S. (2017a) 'Relationship between dental anxiety and pain experience during dental extractions', Asian Journal of Pharmaceutical and Clinical Research, p. 458. doi: 10.22159/ajpcr.2017.v10i3.16518.
- Kumar, S. (2017b) 'The emerging role of botulinum toxin in the treatment of orofacial disorders Literature update', Asian Journal of Pharmaceutical and Clinical Research, p. 21. doi: 10.22159/ajpcr.2017.v10i9.16914.
- Kumar, S. and Rahman, R. (2017) 'Knowledge, Awareness, And Practices Regarding Biomedical Waste Management Among Undergraduate Dental Students', Asian Journal of Pharmaceutical and Clinical Research, p. 341. doi: 10.22159/ajpcr.2017.v10i8.19101.
- Kumar, S. and Sneha, S. (2016) 'Knowledge and awareness regarding antibiotic prophylaxis for infective endocarditis among undergraduate dental students', Asian Journal of Pharmaceutical and Clinical Research, p. 154. doi: 10.22159/ajpcr.2016.v9s2.13405.

- Lekha, L. et al. (2014a) 'Schiff base complexes of rare earth metal ions: Synthesis, characterization and catalytic activity for the oxidation of aniline and substituted anilines', Journal of organometallic chemistry, 753, pp. 72–80.
- Lekha, L. et al. (2014b) 'Synthesis, spectroscopic characterization and antibacterial studies of lanthanide(III) Schiff base complexes containing N, O donor atoms', Journal of Molecular Structure, pp. 307–313. doi: 10.1016/j.molstruc.2013.10.014.
- 36. Marimuthu, M. et al. (2018) 'Canonical Wnt pathway gene expression and their clinical correlation in oral squamous cell carcinoma', Indian journal of dental research: official publication of Indian Society for Dental Research, 29(3), pp. 291–297.
- Menon, S. et al. (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', Colloids and surfaces. B, Biointerfaces, 170, pp. 280–292.
- Narasimman, C. (2018) Application of physics forceps for extraction of mutilated tooth to avoid transalveolar extraction method: A Prospective study. masters. Ragas Dental College and Hospital, Chennai. Available at: http://repositorytnmgrmu.ac.in/9911/ (Accessed: 13 June 2020).
- Natarajan, B., Balakrishnan, G. and Thangavelu, K. (2017) 'Comparison of Efficacy of Amoxicillin versus Ciprofloxacin in Postsurgical Management of Transalveolar Extraction', Journal of pharmacy &bioallied sciences, 9(Suppl 1), pp. S187–S190.
- Neelakantan, P.etal. (2015) 'Antibiofilmactivityofthreeir rigationprotocolsactivatedbyultrasonic, diode laser or Er:YAG laser in vitro', International endodontic journal, 48(6), pp. 602–610.
- Neelakantan, P. et al. (2015) 'Influence of Irrigation Sequence on the Adhesion of Root Canal Sealers to Dentin: A Fourier Transform Infrared Spectroscopy and Push-out Bond Strength Analysis', Journal of endodontia, 41(7), pp. 1108–1111.
- Neelakantan, P., Grotra, D. and Sharma, S. (2013) 'Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis', Journal of endodontia, 39(7), pp. 893–896.
- Ozmen, B. (2019) 'Evaluation of Permanent First Molar Tooth Loss in Young Population from North Turkey', Balkan Journal of Dental Medicine, pp. 20– 23. doi: 10.2478/bjdm-2019-0004.
- Packiri, S., Gurunathan, D. and Selvarasu, K. (2017) 'Management of Paediatric Oral Ranula: A Systematic Review', Journal of clinical and diagnostic research: JCDR, 11(9), pp. ZE06–ZE09.
- Parthasarathy, M. et al. (2016) 'Effect of hydrogen on ethanol-biodiesel blend on performance and emission characteristics of a direct injection diesel engine', Ecotoxicology and environmental safety, 134(Pt 2), pp. 433–439.
- 46. Patil, S. B. et al. (2017) 'Comparison of Extended Nasolabial Flap Versus Buccal Fat Pad Graft in the Surgical Management of Oral Submucous Fibrosis: A Prospective Pilot Study', Journal of maxillofacial and oral surgery, 16(3), pp. 312–321.
- 47. Patturaja, K. and Pradeep, D. (2016) 'Awareness of Basic Dental Procedure among General Population',

Research Journal of Pharmacy and Technology, p. 1349. doi: 10.5958/0974-360x.2016.00258.4.

- Polat, H. B., Ay, S. and Kara, M. I. (2007) 'Maxillary tuberosity fracture associated with first molar extraction: a case report', European journal of dentistry, 1(4), pp. 256–259.
- 49. PradeepKumar, A. R. et al. (2016) 'Diagnosis of Vertical Root Fractures in Restored Endodontically

Treated Teeth: A Time-dependent Retrospective Cohort Study', Journal of endodontia, 42(8), pp. 1175–1180.

 Praveen, K. et al. (2001) 'Hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study', The British journal of oral & maxillofacial surgery, 39(2), pp. 138–140.