

## ORIGINAL ARTICLE

### Assessment of efficacy of balloon tamponade in the management of postpartum haemorrhage

Jyoti Sultania

Associate Professor, Department of Obstetrics & Gynaecology, Major S D Singh Medical College & Hospital, Farukhabad, Uttar Pradesh, India

#### ABSTRACT:

**Background:** Post-partum hemorrhage (PPH) remains a major cause of maternal death with uterine atony implicated in up to 80% of all bleedings. The present study was conducted to assess efficacy of balloon tamponade in the management of postpartum hemorrhage. **Materials & Methods:** 45 patients of postpartum hemorrhage were managed with balloon tamponade. **Results:** The mean height was 63.2 cms, weight was 145.2 kgs, gravida was 2.5 and gestational age was 37.2 weeks. Pre-balloon and post-balloon BMI (Kg/m<sup>2</sup>) was 27.2 and 28.9, multiple pregnancies was 24 and 11, history of uterine surgery was seen in 11 and 14, mean haemoglobin was 12.4 and 12.4 respectively. The difference was significant (P< 0.05). There were 25 patients with any invasive procedure and 12 and overall transfusion was seen in 42 and 16 in pre-balloon and post-balloon cases respectively. The difference was significant (P< 0.05). **Conclusion:** The implementation of IUTB in PPH protocol for caesarean delivery was associated with a drastic drop in second-line invasive procedures.

**Key words:** Caesarean, Post-partum hemorrhage, balloon tamponade

**Corresponding author:** Jyoti Sultania, Associate Professor, Department of Obstetrics & Gynaecology, Major S D Singh Medical College & Hospital, Farukhabad, Uttar Pradesh, India

**This article may be cited as:** Sultania J. Assessment of efficacy of balloon tamponade in the management of postpartum haemorrhage. J Adv Med Dent Scie Res 2016;4(1):209-211.

#### INTRODUCTION

Post-partum hemorrhage (PPH) remains a major cause of maternal death with uterine atony implicated in up to 80% of all bleedings. First-line treatment includes oxytocin followed, when needed, by prostaglandins.<sup>1</sup> In case of failure of these first-line treatments, second-line invasive therapies such as artery ligation, uterine compressive suture and/or radiological artery embolization have to be implemented. Each of these treatments can be performed alone or they may be also combined, with failure leading to hysterectomy. Because of the invasive character and inherent morbidity of these second-line therapies.<sup>2</sup>

Balloon tamponade has become an important part of our armamentarium for the management of Postpartum hemorrhage (PPH).<sup>3</sup> There are now several different devices being used for this purpose, some specifically designed for PPH and some used off-label. These balloon devices have been used following vaginal delivery and caesarean section, in women with postpartum uterine atony and in patients with contracted uteri but bleeding from other causes such as placenta accrete or post abortal hemorrhage.<sup>4</sup> Challenges during balloon filling include the requirement to apply high amounts of physical force to, and repeatedly fill and empty, the syringe used to

deliver the fluid to expand the balloon, which can cause delay in achieving adequate tamponade.<sup>5</sup> The present study was conducted to assess efficacy of balloon tamponade in the management of postpartum hemorrhage.

#### MATERIALS & METHODS

The present study comprised of 45 patients of postpartum hemorrhage. All patients were enrolled after obtaining their written consent.

Data such as name, age etc. was recorded. Obstetric & tamponade system (OTS) was used which had a uterine balloon and a vaginal balloon. Vaginal bleeding can also be controlled by the vaginal tamponade balloon. In all females, blood pressure prior to balloon inflation was recorded. The time to maximum inflation, volume of saline infused, and pressure within the balloon were recorded. All underwent ultrasound evaluation of the uterus before and during the balloon inflation and colour and pulsed wave Doppler examinations of the uterine artery flow velocity pattern were recorded. The amount of blood draining from the drainage channel of the catheter was recorded. Data thus obtained were subjected to statistical analysis. P value < 0.05 was considered significant.

#### RESULTS

**Table I Baseline characteristics**

Parameters	Mean	SD
Height	63.2	2.7
Weight	145.2	7.1
Gravida	2.5	1.1

Gestational age (weeks)	37.2	1.8
-------------------------	------	-----


Table I shows that mean height was 63.2 cms, weight was 145.2 kgs, gravida was 2.5 and gestational age was 37.2 weeks.

**Table II Assessment of parameters**

Parameters	Pre-Balloon	Post-Balloon	P value
BMI (Kg/m <sup>2</sup> )	27.2	28.9	0.12
Multiple pregnancies	24	11	0.01
History of uterine surgery	11	14	0.05
Haemoglobin	12.4	12.4	1

Table II, graph I shows that pre-balloon and post-balloon BMI (Kg/m<sup>2</sup>) was 27.2 and 28.9, multiple pregnancies was 24 and 11, history of uterine surgery was seen in 11 and 14, mean haemoglobin was 12.4 and 12.4 respectively. The difference was significant (P< 0.05).

**Graph I Assessment of parameters**



**Table III Rates of the various invasive procedures implemented**

Outcome	Pre-Balloon	Post-Balloon	P value
Patients with any invasive procedure	25	12	0.01
Overall transfusion	42	16	0.05

Table III shows that there were 25 patients with any invasive procedure and 12 and overall transfusion was seen in 42 and 16 in pre- balloon and post- balloon cases respectively. The difference was significant (P< 0.05).

## DISCUSSION

Post-partum hemorrhage (PPH) remains a major cause of maternal death with uterine atony implicated in up to 80% of all bleedings.<sup>6</sup> First-line treatment includes oxytocin followed, when needed, by prostaglandins.<sup>7</sup> In case of failure of these first-line treatments, second-line invasive therapies such as artery ligation, uterine compressive suture and/or radiological artery embolization have to be implemented. Each of these treatments can be performed alone or they may be also combined, with failure leading to hysterectomy.<sup>8</sup> Because of the invasive character and inherent morbidity of these second-line therapies, alternative approaches have been suggested: the Blakemore probe

first, and then a few other devices, among which the intra-uterine tamponade balloon (IUTB) has triggered much interest.<sup>9</sup> The present study was conducted to assess efficacy of balloon tamponade in the management of postpartum hemorrhage.

We found that the mean height was 63.2 cms, weight was 145.2 kgs, gravida was 2.5 and gestational age was 37.2 weeks. Mishra et al<sup>10</sup> found that the mean heights of patients was 62± 2.7 inches, weight (145± 12 pounds), Gravidity/Parity (2 [1–6]/1), type of delivery was vaginal (12) and caesarean (4) and gestational age was 37.6± 1.4 weeks. The mean systolic blood pressure was 106.42 mm Hg, maximum volume infusion was 960 mL, pressure at maximum

value was 58.22 mm Hg and estimated blood loss was 440 ml.

We found that pre-balloon and post-balloon BMI (Kg/m<sup>2</sup>) was 27.2 and 28.9, multiple pregnancies was 24 and 11, history of uterine surgery was seen in 11 and 14, mean haemoglobin was 12.4 and 12.4 respectively. Soued et al<sup>11</sup> included women with postpartum hemorrhage requiring potent uterotonic treatment with prostaglandins after caesarean delivery over a 9-year period were eligible. The primary outcome was the rate of invasive procedure (conservative surgery, radiological embolization and/or hysterectomy).  $p < 0.05$  was considered statistically significant. A total of 279 patients were included (140 vs. 139). Most baseline characteristics were comparable between the two studied periods. The success rate of the intra-uterine tamponade balloon was 82%, and no related complications occurred. Rates of invasive procedures and transfusion were significantly reduced (28.6% vs. 11.5%,  $p < 0.001$  and 44.3% vs. 28.1%,  $p = 0.006$  respectively) during the “post-balloon” period, and length of hospital stay was shorter ( $p < 0.001$ ). Implementation of intra-uterine tamponade balloon during post-partum hemorrhage after caesarean delivery appears to be safe and effective, with a decrease in both invasive procedures and transfusion rates.

We observed that there were 25 patients with any invasive procedure and 12 and overall transfusion was seen in 42 and 16 in pre- balloon and post- balloon cases respectively. Cho Y et al<sup>12</sup> demonstrated cessation of bleeding in a PPH patient after 320 mL of saline was placed in an intrauterine Senstaken Blakemore tube. Postplacement ultrasound in their case demonstrated that the inflated balloon was not in the fundus but in the lower uterine segment and that it was not tamponading an area of myometrium sufficient to be effective at controlling PPH. They suggested that the mechanism of action was by uterine artery compression rather than direct myometrial tamponade.

The limitation the study is small sample size.

## CONCLUSION

Authors found that the implementation of IUTB in PPH protocol for caesarean delivery was associated with a drastic drop in second-line invasive procedures.

## REFERENCES

1. Bieniarz J, Branda LA, Maqueda E, Morozovsky J, Caldeyro-Barcia R. Aortocaval compression by the uterus in late pregnancy- Unreliability of the sphygmomanometric method in estimating uterine artery pressure. *Am J Obstet Gynecol.* 1968; 102:1106–1115.
2. Smith WC. Uterine tamponade with oxidized gauze in a case of total separation of the placenta with concealed hemorrhage. *N Y State J Med.* 1949; 49: 2187.
3. Georgiou C. Balloon tamponade in the management of postpartum haemorrhage: a review. *BJOG* 2009; 116:748– 757.
4. Bakri YN. Balloon device for control of obstetrical bleeding. *Eur J Obstet Gynecol Reprod Biol* 1999; 86:84.
5. Bakri YN, Amri A, Abdul Jabbar F. Tamponade-balloon for obstetrical bleeding. *Int J Gynaecol Obstet* 2001; 74:139– 142.
6. Belfort MA, Dildy GA, Garrido J, White GL. Intraluminal pressure in a uterine tamponade balloon is curvilinearly related to the volume of fluid infused. *Am J Perinatol* 2011; 8:659-666.
7. Kramer MS, Berg C, Abenhaim H, et al. Incidence, risk factors, and temporal trends in severe postpartum hemorrhage. *Am J Obstet Gynecol* 2013; 5:449-449.
8. Soyama H., Miyamoto M., Ishibashi H., Nakatsuka M., Kawachi H., Sakamoto T., Matuura H., Kuwahara M., Takasaki K., Iwahashi H., et al. Analysis of prophylactic Bakri balloon tamponade failure in patients with placenta previa. *Taiwan J. Obstet. Gynecol.* 2019;58:159–163.
9. Beckmann M.M., Chaplin J. Bakri balloon during cesarean delivery for placenta previa. *Int. J. Gynaecol. Obstet.* 2014;124:118–122.
10. Mishra A. Efficacy of balloon tamponade in the management of postpartum hemorrhage: A clinical study. *International Journal of Clinical Obstetrics and Gynaecology* 2018; 2(1): 07-09.
11. Soued M, Vivanti AJ, Smiljkovski D, Deffieux X, Benachi A, Le Gouez A, Mercier FJ. Efficacy of Intra-Uterine Tamponade Balloon in Post-Partum Hemorrhage after Caesarean Delivery: An Impact Study. *Journal of Clinical Medicine.* 2020 Dec 28;10(1):81.
12. Cho Y, Rizvi C, Uppal T, Condous G. Ultrasonographic visualization of balloon placement for uterine tamponade in massive primary postpartum hemorrhage. *Ultrasound Obstet Gynecol* 2008; 32:711-713.