

## Original Research

### A Clinical Study to evaluate Ocular manifestations in Diabetic Patients

Chander Mohan Kohli<sup>1</sup>, Gaurav Mohan Kohli<sup>2</sup>

<sup>1</sup>Associate Professor and in charge Vitreo Retinal Unit, Hind Institute of Medial Sciences, Safedabad. UP, India;

<sup>2</sup>Consultant Vitreo Retinal Services, Satguru Netrachakitsalaya, Chitrakoot, MP,India

#### **ABSTRACT:**

**Background:** Diabetic retinopathy is the most well-known ocular complication of diabetes and the leading cause of blindness among people 20–64 years of age. Hence; we planned the present study to assess the incidence of different ocular manifestations of diabetes mellitus. **Materials & methods:** A total of 400 diabetic patients were screened during the study period. A registered and experienced ophthalmologist was appointed for ocular screening of all the patients in the present study. Visual acuity of each patient was assessed. Ocular pathologies, if present, were recorded and Microsoft excel sheet and were assessed by SPSS software. Chi- square test was used for assessment of level of significance. **Results:** The overall prevalence of ocular lesions among diabetic patients was 14 percent. Among these ocular lesions, the most common was glaucoma, diabetic retinopathy, cataract and diabetic papillopathy. **Conclusion:** Ocular lesions are prevalent in significant proportion in diabetic patients. Effective glycemic control has been demonstrated to reduce both the incidence and progression of DR.

**Key words:** Diabetes, ocular, retinopathy.

Received: 10 June, 2019

Revised: 12 July 2019

Accepted: 14 July 2019

Corresponding Author: Dr. Gaurav Mohan Kohli, Consultant Vitreo Retinal Services, Satguru Netrachakitsalaya, Chitrakoot, MP, India

This article may be cited as: Kohli CM, Kohli GM. A Clinical Study to evaluate Ocular manifestations in Diabetic Patients. J Adv Med Dent Scie Res 2019;7(8):247-250.

#### **INTRODUCTION**

Complications of diabetes mellitus (DM) are progressive and almost resulting by chronic exposure to high blood levels of glucose caused by impairments in insulin metabolism and biological macromolecules such as carbohydrates, lipids, proteins and nucleic acids. DM and its complications are rapidly becoming the world's most significant cause of morbidity and mortality<sup>1-3</sup> It can affect children, young people and adults and is becoming more common. Cataract is a major cause of vision impairment in people with diabetes. Numerous studies have documented an association between diabetes and cataracts.<sup>4,5</sup> This association is supported by an abundance of data from clinical epidemiological studies and basic science studies.<sup>6-8</sup> Hence; we planned the present study to assess the incidence of different ocular manifestations of diabetes mellitus.

#### **MATERIALS & METHODS**

The present study was conducted to assess the incidence of different ocular manifestations of diabetes mellitus. A total of 460 diabetic patients were screened during the study period. Subjects with history of any other systemic illness, known drug allergy, any other co-morbid condition, malignancy of ocular region were excluded from study group.

After meeting the exclusion criteria, a total of 60 subjects were excluded from the present study. So the net sample size was reduced to 400.

Patient's basic information was noted first, which includes name, age, sex, reg. no, occupation etc. patient's postal address was also noted in detail in order to call them again in future for follow up if required.

Fasting blood samples were obtained from all the patients for confirming the diagnosis of diabetes. A registered and experienced ophthalmologist was appointed for ocular

screening of all the patients in the present study. Visual acuity of each patient was assessed with a Snellen chart, and pinhole visual acuity was used to screen for refractive errors. Ocular pathologies, if present, were recorded and Microsoft excel sheet and were assessed by SPSS software. Chi- square test was used for assessment of level of significance.

A total of 400 diabetic patients were screened in the present study. Among these diabetic patients, 240 were males, while the remaining 160 were females. 25 percent of the patients were less than 30 years of age, 50 percent of the patients belonged to the age group of 30 to 50 years and the remaining 25 percent of the patients were more than 50 years of age. The overall prevalence of ocular lesions among diabetic patients was 14 percent. Among these ocular lesions, the most common was glaucoma, diabetic retinopathy, cataract and diabetic papillopathy.

**RESULTS**

**Table 1:** Distribution of subjects Age-wise and gender-wise

Parameter		Number of subjects	Percentage
Age-group (years)	Less than 30	100	25
	30 to 50	200	50
	More than 50	100	25
Gender	Male	240	60
	Female	160	40

**Table 2:** Prevalence of ocular lesions among diabetic

S No.	Ocular lesion	Number of subjects	Percentage
1	Diabetic retinopathy	12	3
2	Diabetic Papillopathy	10	2.5
3	Glaucoma	14	3.5
4	Cataract	12	3
5	Others	8	2
Total		56	14

**DISCUSSION**

DM as a systemic disease has several well-known ocular complications including anterior and posterior segment such as DES, glaucoma, corneal pathologies, and retinopathy. In the current study, we checked the metabolic status of Type 1 DM children with a full anterior and posterior segment ophthalmologic examination and compared the results with sex- and age-matched healthy controls.

In the present study, a total of 400 diabetic patients were screened in the present study. Among these diabetic patients, 240 were males, while the remaining 160 were females. 25 percent of the patients were less than 30 years of age, 50 percent of the patients belonged to the age group of 30 to 50 years and the remaining 25 percent of the patients were more than 50 years of age. The overall prevalence of ocular lesions among diabetic patients was 14 percent. Among these ocular lesions, the most common was glaucoma, diabetic retinopathy, cataract and diabetic papillopathy.

Cairncross JP et al<sup>9</sup> determined the prevalence of eye pathology in a group of diabetic patients at National District Hospital by screening for diabetes-associated ocular pathology. Interviews were used to collect information

regarding diabetic patients’ history of diabetes mellitus and if and when previous diabetic retinopathy screening was performed. Visual acuity was assessed, intra-ocular pressure measured and a non-mydriatic digital fundus camera used to screen for retinal pathology. During the last year, only 4.5% of patients had their vision checked with a Snellen chart, and 16.5% were examined with an ophthalmoscope. Since diagnosis of diabetes, only 15.5% of patients were referred to an ophthalmologist. Patient referral was needed for 87 (42.9%) cases for refractive disorders, 37 (18.2%) for suspected glaucoma, 30 (14.8%) for cataracts, and 22 (10.8%) for diabetic retinopathy. This study confirmed that glaucoma, cataracts and diabetic retinopathy are prevalent eye conditions among diabetic patients. Offering eye screening at primary healthcare level may contribute to early detection of eye pathology and timeous referral for sight-saving treatment.<sup>9</sup>

In the present study, the overall prevalence of ocular lesions among diabetic patients was 14 percent. Among these ocular lesions, the most common was glaucoma, diabetic retinopathy, cataract and diabetic papillopathy. Webb EM et al<sup>10</sup> determined the prevalence of diabetic retinopathy, maculopathy and visual loss in primary care patients and to

identify associated risk factors. They conducted a cluster randomised trial at primary care clinics in the Tshwane district in South Africa. Grades of retinopathy and maculopathy (with fundus camera) and visual acuity (Snellen chart) were assessed and, using mobile screening and teleophthalmology, clinical and biochemical testing was conducted to obtain information about glycaemic control and microvascular complications. The prevalence rates for any retinopathy, preproliferative retinopathy and proliferative retinopathy were 24.9, 19.5 and 5.5%, respectively. The prevalence rates of diabetic maculopathy, observable maculopathy and referable maculopathy were 20.8, 11.8 and 9.0%, respectively. The presence of retinopathy was associated with high body mass index, systolic blood pressure, being on insulin treatment, high HbA1c and the presence of neuropathy. High systolic blood pressure, being on insulin treatment, high HbA1c level and high low-density lipoprotein cholesterol level as well as the presence of albuminuria were significant in predicting any diabetic maculopathy. Laser photocoagulation was given to 8.3% of patients from the mobile unit and 12% of patients were referred to the nearest hospital with an outpatient eye clinic for follow-up treatment of various other eye conditions. Using the WHO categories, the study found that 78.1% of diabetes patients had normal vision, 19.3% were visually impaired and 2.2% were severely impaired or blind. High prevalence rates for diabetic retinopathy, maculopathy and visual loss were found and associations were identified.<sup>10</sup> Lawan A et al<sup>11</sup> determined the pattern of retinopathy seen in diabetic patients attending the outpatient clinic in Aminu Kano Teaching Hospital, Kano, Nigeria. Consecutive patients who were attending the diabetic clinic and who consented were examined over a three-month period. Information obtained includes patient's bio data, type and duration of disease, and findings on eye examination. The fundus was examined with direct and indirect ophthalmoscopes, +90 D with slit lamp and fundal photography. Retinopathy was graded using the International Clinical Diabetic Retinopathy Disease Severity Scale (ICDRDSS). A total of 214 patients were examined during the study period. There were 88 males and 126 females (M: F = 1: 1.43). The mean age of the study population was  $52.14 \pm 13.23$  years. The mean age of patients without diabetic retinopathy (DR) was  $49.14 \pm 13.17$  years and the mean age of patients with DR was  $58.51 \pm 10.94$  years. Forty nine patients (23%) had insulin-dependent diabetes mellitus (IDDM) while 165 patients (77%) had non insulin dependent diabetes mellitus (NIDDM). There was statistically significant difference in presence of retinopathy in patients with IDDM compared to those with NIDDM [ $X(2) = 29.77$  {95% CI},  $P = 0.000$ ]. DR was significantly more common in patients with disease duration of 15 years or more compared with those with disease duration of 14 years or less [ $X(2) = 65.85$ , {95% CI}  $P = 0.000$ ]. Based on ICDRDSS scale, 136 patients (64%) had no retinopathy and 78 patients (36%) had

retinopathy. Some patients were visually impaired and the cause of blindness was DR in 6 patients (2.8%). Cataract and glaucoma were the cause in 6 patients (2.8%). Diabetic retinopathy is common in our environment and is more frequent in IDDM and those with long disease duration.<sup>11</sup> Although specific treatment modalities for retinopathy threatening vision have improved over years of clinical and research experience, importance of preventive measures (tight glycaemic and blood pressure (BP) control, smoking cessation, regular eye screening) cannot be underestimated. Routine comprehensive eye evaluation helps to detect early treatable stages of DR which are frequently asymptomatic. Comprehensive eye examination is recommended within 3 to 5 years of disease onset in type 1 diabetic patients aged 10 years or older and immediately after diagnosis in type 2 diabetes. Diabetic patients without evidence of retinopathy should undergo eye examinations every year to detect its emergence. For patients with moderate-to-severe preproliferative disease, more frequent eye examinations are necessary to determine when to initiate therapy. Kathiara et al<sup>12</sup> also observed cataract to be the most common ocular manifestation of diabetes mellitus 35-patients (58.33%). The second most common manifestation was diabetic retinopathy-25 patients (41.66%). Also the most common associated systemic disease was found to be hypertension-20 patients (33.3%). It is imperative to diagnose these conditions early enough, to reduce the overall ocular morbidity caused by diabetes. It is also very important to closely follow these patients, as various ocular manifestations may develop at any time during their life span and early diagnosis and treatment are of utmost importance.

## CONCLUSION

From the above obtained results; the authors conclude that ocular lesions are prevalent in significant proportion in diabetic patients. Therefore, periodic ocular screening of the diabetic patients should be done to avoid further complications. Effective glycaemic control has been demonstrated to reduce both the incidence and progression of DR.

## REFERENCES

1. Writing Team for the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Research Group. Effect of intensive therapy on the microvascular complications of type 1 diabetes mellitus. *JAMA* 2002;287:2563-9.
2. Forbes JM, Soldatos G, Thomas MC. Below the radar: advanced glycation end products that detour "around the side". Is HbA1c not an accurate enough predictor of long term progression and glycaemic control in diabetes? *ClinBiochem Rev.* 2005;26:123-134.
3. Kernell A, Dedorsson I, Johansson B, Wickström CP, Ludvigsson J, Tuvemo T, *et al.* Prevalence of diabetic retinopathy in children and adolescents with IDDM. *A*

- population-based multicentre study. *Diabetologia* 1997;40:307-10.
4. Singh PP, Mahadi F, Roy A, Sharma P. Reactive oxygen species, reactive nitrogen species and antioxidants in etiopathogenesis of diabetes mellitus type-2. *Indian J ClinBiochem.* 2009;24:324–342.
  5. Moss SE, Klein R, Klein BE. The 14-year incidence of visual loss in a diabetic population. *Ophthalmology.* 1998;105:998–1003.
  6. Kumari S, Panda S, Mangaraj M, Mandal MK, Mahapatra PC. Plasma MDA and antioxidant vitamins in diabetic retinopathy. *Indian J ClinBiochem.* 2008;23:158–162.
  7. Retinopathy and nephropathy in patients with type 1 diabetes four years after a trial of intensive therapy. The Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Research Group. *N Engl J Med.* 2000;342:381–389.
  8. Stratton IM, Kohner EM, Aldington SJ, Turner RC, Holman RR, Manley SE, Matthews DR. UKPDS 50: risk factors for incidence and progression of retinopathy in Type II diabetes over 6 years from diagnosis. *Diabetologia.* 2001;44:156–163.
  9. Cairncross JP, Steinberg WJ, Labuschagne MJ. Prevalence of eye pathology in a group of diabetic patients at National District Hospital Outpatient Department in Bloemfontein, South Africa. *Afr J Prim Health Care Fam Med.* 2017;9(1):e1-e7. Published 2017 Sep 27. doi:10.4102/phcfm.v9i1.1440
  10. Webb EM1, Rheeder P, Roux P. Screening in Primary Care for Diabetic Retinopathy, Maculopathy and Visual Loss in South Africa. *Ophthalmologica.* 2016;235(3):141-9.
  11. Lawan A, Mohammed TB. Pattern of diabetic retinopathy in Kano, Nigeria. *Ann Afr Med.* 2012 Apr-Jun;11(2):75-9. doi: 10.4103/1596-3519.93528.
  12. Kathiara A, Patel R, Ahir H. A Study on Ocular Manifestations of Diabetes Mellitus. *Journal of Research in Medical and Dental Science* 2015;3:143-147.